



09 509098 2CT  
526 Rec'd PCT/PTO 22 MAR 2000

FORM PTO-1390 (Modified) (REV 5-93)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER 053466/0274	
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371					
		U.S. APPLICATION NO. (If known, see 37 C.F.R. 1.5)		UNASSIGNED	
INTERNATIONAL APPLICATION NO. PCT/JP98/04469		INTERNATIONAL FILING DATE 02 OCTOBER 1998		PRIORITY DATE CLAIMED 03 OCTOBER 1997	
TITLE OF INVENTION NATURAL HUMANIZED ANTIBODY					
APPLICANT(S) FOR DO/EO/US Masayuki TSUCHIYA					
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:					
1.	<input checked="" type="checkbox"/>	This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.			
2.	<input type="checkbox"/>	This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371.			
3.	<input type="checkbox"/>	This express request to begin national examination procedures (35 U.S.C. 371(f)) at any time rather than delay examination until the expiration of the applicable time limit set in 35 U.S.C. 371(b) and PCT Articles 22 and 39(1).			
4.	<input checked="" type="checkbox"/>	A proper Demand for International Preliminary Examination was made by the 19 <sup>th</sup> month from the earliest claimed priority date.			
5.	<input checked="" type="checkbox"/>	A copy of the International Application as filed (35 U.S.C. 371(c)(2))			
	<input type="checkbox"/>	is transmitted herewith (required only if not transmitted by the International Bureau).			
	<input checked="" type="checkbox"/>	has been transmitted by the International Bureau.			
	<input type="checkbox"/>	is not required, as the application was filed in the United States Receiving Office (RO/US)			
6.	<input checked="" type="checkbox"/>	A translation of the International Application into English (35 U.S.C. 371(c)(2)).			
7.	<input checked="" type="checkbox"/>	Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))			
	<input type="checkbox"/>	are transmitted herewith (required only if not transmitted by the International Bureau).			
	<input type="checkbox"/>	have been transmitted by the International Bureau.			
	<input type="checkbox"/>	have not been made; however, the time limit for making such amendments has NOT expired.			
	<input checked="" type="checkbox"/>	have not been made and will not be made.			
8.	<input type="checkbox"/>	A translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371(c)(3)).			
9.	<input checked="" type="checkbox"/>	An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).			
10.	<input type="checkbox"/>	A translation of the annexes to the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).			
Items 11. to 16. below concern other document(s) or information included:					
11.	<input checked="" type="checkbox"/>	An Information Disclosure Statement under 37 CFR 1.97 and 1.98.			
12.	<input checked="" type="checkbox"/>	An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.			
13.	<input checked="" type="checkbox"/>	A FIRST preliminary amendment.			
	<input type="checkbox"/>	A SECOND or SUBSEQUENT preliminary amendment.			
14.	<input type="checkbox"/>	A substitute specification.			
15.	<input type="checkbox"/>	A change of power of attorney and/or address letter.			
16.	<input checked="" type="checkbox"/>	Other items or information: Sequence Listing (71 pages); Microorganism Deposit Receipts (12 pages)			



422 Rec'd PCT/PTO 22 MAR 2000

U.S. APPLICATION NO. (If known, see 37 C.F.R. 1.50)		INTERNATIONAL APPLICATION NO.		ATTORNEY'S DOCKET NUMBER	
UNASSIGNED		PCT/JP98/04469		053466/0274	
17. <input checked="" type="checkbox"/> The following fees are submitted:				CALCULATION	
Basic National Fee (37 CFR 1.492(a)(1)-(5): Search Report has been prepared by the EPO or JPO ..... \$840.00					
International preliminary examination fee paid to USPTO (37 CFR 1.482) ..... \$670.00					
No international preliminary examination fee paid to USPTO (37 CFR 1.482) but international search fee paid to USPTO (37 CFR 1.445(a)(2)) ..... \$690.00					
Neither international preliminary examination fee (37 CFR 1.482) nor International search fee (37 CFR 1.445(a)(2)) paid to USPTO ..... \$970.00					
International preliminary examination fee paid to USPTO (37 CFR 1.482) and all claims satisfied provisions of PCT Article 33(2)-(4) ..... \$96.00					
ENTER APPROPRIATE BASIC FEE AMOUNT =				\$840.00	
Surcharge of \$130.00 for furnishing the oath or declaration later than 20 Months from the earliest claimed priority date (37 CFR 1.492(e))					
Claims	Number Filed	Included in Basic Fee	Extra Claims	Rate	
Total Claims	13	-	20	= 0	\$18.0
Independent Claims	3	-	3	= 0	\$78.0
Multiple dependent claim(s) (if applicable)					\$260.00
TOTAL OF ABOVE CALCULATIONS =					\$840.00
Reduction by 1/2 for filing by small entity, if applicable. Verified Small Entity statement must also be filed. (Note 37 CFR 1.9, 1.27, 1.28).					\$0.00
SUBTOTAL =					\$840.00
Processing fee of \$130.00 for furnishing English translation later the 20 months from the earliest claimed priority date (37 CFR 1.492(f)).				+	
TOTAL NATIONAL FEE =					\$840.00
Fee for recording the enclosed assignment (37 CFR 1.21(h)). The assignment must be accompanied by an appropriate cover sheet (37 CFR 3.28, 3.31). \$40.00 per property				+	\$40.00
TOTAL FEES ENCLOSED =					\$880.00
				Amount to be: refunded	\$
				charged	\$
<p>a. <input checked="" type="checkbox"/> A check in the amount of \$880.00 to cover the above fees is enclosed.</p> <p>b. <input type="checkbox"/> Please charge my Deposit Account No. <u>19-0741</u> in the amount of \$880.00 to the above fees. A duplicate copy of this sheet is enclosed.</p> <p>c. <input checked="" type="checkbox"/> The Commissioner is hereby authorized to charge any additional fees which may be required, or credit any overpayment to Deposit Account No. <u>19-0741</u>. A duplicate copy of this sheet is enclosed.</p>					
<p>NOTE: Where an appropriate time limit under 37 CFR 1.494 or 1.495 has not been met, a petition to revive (37 CFR 1.137(a) or (b)) must be filed and granted to restore the application to pending status.</p>					
SEND ALL CORRESPONDENCE TO:					
Foley & Lardner Washington Harbour 3000 K Street, N.W., Suite 500 Washington, D.C. 20007-5109			 SIGNATURE Reg. No. 33,715 HAROLD C. WEGNER REGISTRATION NUMBER 25,258		

422 Rec'd PCT/PTO 22 MAR 2000

## IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

In re patent application of Masayuki TSUCHIYA

Serial No. UNASSIGNED

Filed: March 22, 2000

For: NATURAL HUMANIZED ANTIBODY

## PRELIMINARY AMENDMENT

Assistant Commissioner of Patents  
Washington, D.C. 20231

Sir:

Prior to examination of the above-identified application, Applicants respectfully request that the following amendments be entered into the application:

IN THE CLAIMS:

- |                         |   |
|-------------------------|---|
| Claim 3, line 2,        | delete "or 2".  |
| Claim 5, lines 1 and 2, | delete "any of claims 1 to 4" and insert --claim 1--. |
| Claim 6, lines 2 and 3, | delete "any of claims 1 to 5" and insert --claim 1--. |
| Claim 9, line 2,        | delete "any of claims 6 to 8" and insert --claim 6--. |

REMARKS

Applicants respectfully request that the foregoing amendments to Claims 2, 5, 6 and 9 be entered. These amendments are being made in order to avoid this application incurring a surcharge for the presence of one or more multiple dependent claims. No new matter has been added.

Respectfully submitted,

 Reg. no. 33,715

March 22, 2000  
Date

for Harold C. Wegner  
Reg. No. 25,258

FOLEY & LARDNER  
Washington Harbour  
3000 K Street, N.W., Suite 500  
Washington, DC 20007-5109

422 Rec'd PCT/PTO 22 MAR 2000  
SPECIFICATIONNATURAL HUMANIZED ANTIBODY

## 5 Technical Field

The present invention relates to a method of preparing natural humanized antibody and the natural humanized antibody obtained by said method of preparation. The present invention also relates to DNA encoding natural humanized antibody, an expression vector comprising said DNA, a host comprising said DNA, and a method of preparing natural humanized antibody from cells into which said DNA has been introduced.

## 15 Background Art

Mouse monoclonal antibodies can be relatively easily isolated by the widely used hybridoma technology (Kohler, G. and Milstein, C. Nature (1975) 256, 495-497). On the other hand, a similar technique for human hybridoma has yet to be widespread though it is expected to become so. Furthermore, there is a need for antibodies to human antigens in clinical applications, and therefore the generation of mouse monoclonal antibodies is indispensable for the development of antibody pharmaceuticals.

In fact, a number of monoclonal antibodies have been isolated against tumor cells and viruses, and have been studied in clinical applications. It has been revealed, however, that mouse antibodies, which are a foreign substances to humans, induce HAMA (human anti-mouse antibody) due to the potent antigenicity, and that it is extremely unsuitable for clinical applications because of such problems as a weak activity of inducing ADCC (Schroff, R. W., Cancer Res. (1985) 45, 879-885; Shawler, D. L., et al, J. Immunol. (1985) 135, 1530-1535).

In order to solve this problem, chimeric antibody was created (Neuberger, M. S. et al., Nature (1984) 312,

604-608; Boulianne, G. L. et al., Nature (1984) 312, 643-646). Chimeric antibody is made by linking a variable region of a mouse antibody to a constant region of a human antibody, i.e. in chimeric antibody the constant region of the mouse antibody which is responsible for a particularly potent antigenicity has been replaced with a human counterpart. This is expected to enable a physiological binding with a human Fc receptor and to induce Fc-mediated functions. In fact, marked decreases in antigenicity has been reported in a clinical study using chimeric antibodies (LoBuglio, A. F. et al., Proc. Natl. Acad. Sci. U.S.A. (1989) 86, 4220-4224). However, trouble-causing cases were reported that developed HAMA against mouse variable regions (LoBuglio, A. F. et al., Proc. Natl. Acad. Sci. U.S.A. (1989) 86, 4220-4224).

Accordingly, methods have been developed, though more complicated, for making a humanized antibody which is closer to a human antibody. This is a technique of reconstructing the antigen binding site of a mouse antibody on a human antibody (Jones, P. T. et al., Nature (1986) 321, 5225-525; Verhoeyen, M. et al., Science (1988) 239, 1534-1536; Riechmann, L. et al., Nature (1988) 332, 323-327)). Thus, a variable region of an antibody, for both of the H chain and the L chain, comprises four framework regions (FRs) and three complementarity determining regions (CDRs) sandwiched between them.

It is known that CDR is mainly responsible for the formation of antigen binding sites and some amino acid residues on the FR are involved therein either directly or indirectly. Since the basic structures of antibodies are similar to each other, it was thought possible to graft an antigen binding site of an antibody to another antibody. The research group led by G. Winter has, in fact, successfully grafted CDRs of a mouse anti-rhizobium antibody to a human antibody (CDR-grafting) thereby obtaining a humanized antibody having a rhizobium binding

activity (Jones, P. T. et al., Nature (1986) 321, 522-525).

5 In some cases, however, humanization by CDR-grafting alone does not provide humanized antibody that has an antigen binding activity similar to the original mouse antibody. Accordingly, as described above, attempts have been made to replace some FR amino acid residues. FR amino acid residues to be replaced are involved in the maintenance of the structure of amino acid residues that  
10 constitute the basic structure of an antibody molecule (canonical structure; Chothia, C. et al., Nature (1989) 342, 877-883; Chothia, C. and Lesk, A. M. J. Molec. Biol. (1987) 196, 901-917) or CDRs, or directly interact with antigen molecules.

15 In fact, amino acid substitution on the FR has been made for most of the humanized antibody, wherein artificial FR sequences that do not naturally occur are formed. At times, too many amino acid substitutions have been made, which makes doubtful the original meaning of  
20 CDR-grafting for minimizing the antigenicity of mouse antibody (Queen, C. et al., Proc. Natl. Acad. Sci. U.S.A. (1989) 86, 10029-10033; Co, M. S. et al., Proc. Natl. Acad. Sci. U.S.A. (1991) 88, 2869-2873).

25 A solution to this problem is to devise methods of selecting human FRs. Thus, the number of FR amino acid residues to be replaced depends on the homology between the FRs of the human antibody selected for CDR-grafting and the FRs of the original mouse antibody. Accordingly, human FRs having a high homology with mouse FRs are  
30 usually selected so as to minimize the degree of substitution. However, in many cases even the FRs of humanized antibody thus obtained have amino acid sequences that do not occur naturally, which may present the problem of antigenicity. Thus, there is a need for  
35 the technology of constructing humanized antibody that can solve the above problems, have lower probability of inducing antigenicity, and have higher safety.

# Disclosure of the Invention

5 The present invention is an improvement of the conventional method of constructing humanized antibody, and provides a method of constructing humanized antibody that completely retains the antigen binding activity of the original mouse antibody and that comprises naturally occurring human FRs, in other words a method of constructing humanized antibody that involves no amino acid substitution on the FR.

10 Thus, the present invention provides a method of preparing a natural humanized antibody which comprises conducting a homology search for the FR of a primary design antibody and selecting a natural human FR retaining the artificial amino acid residues contained in the FR of the primary design antibody and having a homology therewith. As used herein, the primary design antibody is a humanized antibody (also called a reshaped human antibody) prepared by the conventional CDR-grafting.

15 The present invention also provides a method of preparing a natural humanized antibody which comprises conducting a homology search for the FR of a primary design antibody, selecting a natural human FR retaining the artificial amino acid residues contained in the FR of the primary design antibody and having a homology therewith, and exchanging one or a plurality of different amino acid residues between the FR of the primary design antibody and the selected natural human FR.

20 Preferably, in the above method of preparation, the primary design antibody comprises the CDRs derived from a first animal species and the FRs derived from a second animal species. More preferably, in the primary design antibody the first animal species is a non-human mammal and the second animal species is human. Examples of the first animal species, i.e. a mammal, include mouse, rat, hamster, rabbit, and monkey.



002220 86060550

The present invention also provides a method of preparing a natural humanized antibody which comprises conducting a homology search for the FR of a primary design antibody, selecting a natural human FR retaining the artificial amino acid residues derived from the FR of a non-human antibody contained in the FR of the primary design antibody and having a high homology therewith, and exchanging one or a plurality of different amino acid residues between the FR of the primary design antibody and the selected natural human FR.

The present invention also provides a natural humanized antibody obtained by the above preparation method.

The present invention also provides a natural humanized antibody containing the CDRs derived from a first animal species and the FRs derived from a second animal species characterized in that said FRs comprise an amino acid sequence which is different from the FRs used for CDR-grafting by one or a plurality of amino acid residues and is replaced with the FR derived from the second animal species having the same amino acid residues as said different amino acid residues at the same positions. Preferably the first animal species is a non-human mammal and the second animal species is human. Examples of the first animal species, i.e. a mammal, include mouse, rat, hamster, rabbit, and monkey.

The present invention also provides DNA encoding the above natural humanized antibody.

The present invention also provides an expression vector comprising the above DNA.

The present invention also provides a host comprising the above DNA.

The present invention also provides a method of preparing a natural humanized antibody which comprises culturing cells into which an expression vector comprising the above DNA has been introduced and collecting the desired natural humanized antibody from

the culture of said cells.

The present invention also provides a pharmaceutical composition comprising a natural humanized antibody.

5      Brief Explanation of the Drawings

Figure 1 is a graph showing that the fluorescent intensity of chimeric anti-HM1.24 antibody is shifted similarly to that of mouse anti-HM1.24 antibody as compared to control antibody in the FCM analysis using a human myeloma cell line KPMM2.

Figure 2 is a graph showing that chimeric anti-HM1.24 antibody inhibits the binding of biotinylated mouse anti-HM1.24 antibody to the WISH cells in a dose-dependent manner similarly to that of mouse anti-HM1.24 antibody.

Figure 3 is a graph showing that chimeric anti-HM1.24 antibody has an increased cytotoxic activity to the RPMI 8226 cells with increasing E/T ratios whereas control IgG1 or mouse anti-HM1.24 antibody has no cytotoxic activity to the RPMI 8226 cells.

Figure 4 is a diagram showing a method of constructing the L chain of reshaped human anti-HM1.24 antibody by CDR-grafting using the PCR method.

Figure 5 is a diagram showing a method of constructing the H chain of reshaped human anti-HM1.24 antibody in which oligonucleotides RVH1, RVH2, RVH3, and RVH4 are assembled by the PCR method.

Figure 6 is a diagram showing a method of constructing the H chain V region of human-mouse hybrid anti-HM1.24 antibody.

Figure 7 is a diagram showing a method of constructing the H chain V region of mouse-human hybrid anti-HM1.24 antibody.

Figure 8 is a graph showing that the L chain version a of reshaped human anti-HM1.24 antibody has an antigen binding activity of a similar degree to that of chimeric anti-HM1.24 antibody. In the figure, -1 and -2 represent

different lots.

Figure 9 is a graph showing the antigen binding activity of reshaped human anti-HM1.24 antibody prepared from a combination of the L chain version a and the H chain version a, b, f, or h, and chimeric anti-HM1.24 antibody.

Figure 10 is a graph showing the antigen binding activity of reshaped human anti-HM1.24 antibody prepared from a combination of the L chain version b and the H chain version a, b, f, or h, and chimeric anti-HM1.24 antibody.

Figure 11 is a graph showing the binding inhibition activity of reshaped human anti-HM1.24 antibody prepared from a combination of the L chain version a and the H chain version a, b, f, or h, and chimeric anti-HM1.24 antibody.

Figure 12 is a graph showing the binding inhibition activity of reshaped human anti-HM1.24 antibody prepared from a combination of the L chain version b and the H chain version a, b, f, or h, and chimeric anti-HM1.24 antibody.

Figure 13 is a graph showing the antigen binding activity of the H chain versions a, b, c, and d of reshaped human anti-HM1.24 antibody and chimeric anti-HM1.24 antibody.

Figure 14 is a graph showing the antigen binding activity of the H chain versions a and e of reshaped human anti-HM1.24 antibody and chimeric anti-HM1.24 antibody. In the figure, -1 and -2 represent different lots.

Figure 15 is a graph showing the binding inhibition activity of the H chain versions a, c, p, and r of reshaped human anti-HM1.24 antibody and chimeric anti-HM1.24 antibody.

Figure 16 is a graph showing the antigen binding activity of human-mouse hybrid anti-HM1.24 antibody, mouse-human hybrid anti-HM1.24 antibody and chimeric

anti-HM1.24 antibody.

Figure 17 is a graph showing the antigen binding activity of the H chain version a, b, c, and f of reshaped human anti-HM1.24 antibody and chimeric anti-HM1.24 antibody.

Figure 18 is a graph showing the antigen binding activity of the H chain versions a and g of reshaped human anti-HM1.24 antibody and chimeric anti-HM1.24 antibody.

Figure 19 is a graph showing the binding inhibition activity of the H chain versions a and g of reshaped human anti-HM1.24 antibody and chimeric anti-HM1.24 antibody.

Figure 20 is a graph showing the antigen binding activity of the H chain versions h and i of reshaped human anti-HM1.24 antibody and chimeric anti-HM1.24 antibody.

Figure 21 is a graph showing the antigen binding activity of the H chain versions f, h, and j of reshaped human anti-HM1.24 antibody and chimeric anti-HM1.24 antibody.

Figure 22 is a graph showing the binding inhibition activity of the H chain versions h and i of reshaped human anti-HM1.24 antibody and chimeric anti-HM1.24 antibody.

Figure 23 is a graph showing the binding inhibition activity of the H chain versions f, h, and j of reshaped human anti-HM1.24 antibody and chimeric anti-HM1.24 antibody.

Figure 24 is a graph showing the antigen binding activity of the H chain versions h, k, l, m, n, and o of reshaped human anti-HM1.24 antibody and chimeric anti-HM1.24 antibody.

Figure 25 is a graph showing the antigen binding activity of the H chain versions a, h, p, and q of reshaped human anti-HM1.24 antibody and chimeric anti-HM1.24 antibody.

Figure 26 is a graph showing the binding inhibition activity of the H chain versions h, k, l, m, n, and o of reshaped human anti-HM1.24 antibody and chimeric anti-HM1.24 antibody to the WISH cells.

5        Figure 27 is a graph showing the binding inhibition activity of the H chain versions a, h, p, and q of reshaped human anti-HM1.24 antibody and chimeric anti-HM1.24 antibody.

10       Figure 28 is a graph showing the antigen binding activity of the H chain versions a, c, p, and r of reshaped human anti-HM1.24 antibody and chimeric anti-HM1.24 antibody.

15       Figure 29 is a graph showing that natural humanized anti-HM1.24 antibody (the secondary design antibody) has an antigen binding activity of a similar degree to that of reshaped human anti-HM1.24 antibody (the primary design antibody).

20       Figure 30 is a graph showing that natural humanized anti-HM1.24 antibody (the secondary design antibody) has a binding inhibition activity of a similar degree to that of reshaped human anti-HM1.24 antibody (the primary design antibody).

25       Figure 31 is a graph showing that purified reshaped human anti-HM1.24 antibody has an antigen binding activity of a similar degree to that of chimeric human anti-HM1.24 antibody.

30       Figure 32 is a graph showing that purified reshaped human anti-HM1.24 antibody has an binding inhibition activity of a similar degree to that of chimeric human anti-HM1.24 antibody.

35       Figure 33 is a graph showing that natural humanized anti-HM1.24 antibody (the secondary design antibody) has an increased cytotoxic activity to the KPMM2 cells with increasing E/T ratios.

Embodiment for Carrying Out the Invention  
1. Natural FR sequence

In order to produce antibodies to a variety of antigens from the genes comprising limited antibody variable regions, organisms have a mechanism of introducing random gene mutations (called somatic mutations) in the antibody variable regions. In theory this should form extremely diverse FR amino acid sequences, but in practice positions of amino acid residues more prone to the introduction of mutations and the kinds of amino acid residues appear to be limited to a certain degree as determined by structural analysis of many human antibody FRs for which actual structures have been elucidated.

As used herein, the term FR refers to the FR that has been defined in Kabat, E. A. et al., Sequence of Proteins of Immunological Interest (1991). Thus, in the H chain, FR1 is amino acids No. 1 to 30, FR2 is amino acids No. 36 to 49, FR3 is amino acids No. 66 to 94, and FR4 is amino acids No. 103 to 113. On the other hand, in the L chain FR1 is amino acids No. 1 to 23, FR2 is amino acids No. 35 to 49, FR3 is amino acids No. 57 to 88, and FR4 is amino acids No. 98 to 107.

## 2. From human FR to natural human FR

In many cases, humanized antibodies (also called reshaped human antibody) produced by the conventional CDR-grafting method have FR amino acid sequences that cannot be found in nature. However, because a variety of FR amino acid sequences have already been found by somatic mutation as mentioned above, it is possible that FRs having artificial amino acid residues created by humanization could be converted into human FRs that occur in nature.

The present invention is intended to create humanized antibody comprising naturally occurring human FRs in stead of artificial FRs by further processing humanized antibody that was constructed by the conventional humanization technology. When humanized antibody that underwent amino acid substitution is

subjected to homology search using human antibody FRs and known databases such as Swiss Plot (protein sequence database), GenBank (nucleic acid sequence database), PRF (protein sequence database) PIR (protein sequence database), and GenPept (translanted protein sequence from GenBank), human FRs having completely matched amino acid sequences or human FRs having homology can be found.

In the former case, FR substitution was carried out when seen from the human FR that was used as the acceptor of CDR-grafting, in which a formed FR that had been presumed to be artificial is present in the natural FR, which can be considered an acceptor, and therefore an FR that underwent no FR substitution can be obtained. In the latter case, by focusing on the amino acid sequence of human FR having a high homology with an artificial FR, it is possible to effect amino acid substitution in the artificial FR that results in returning to a suitable natural human antibody thereby causing a complete match with the natural human FR. This procedure represents humanization on CDR-grafted antibodies.

Since homology search of amino acid sequences between human antibodies is conducted in this case, it is possible to find a human FR that belongs to the same subgroup as the human FR used in CDR-grafting and to find an amino acid sequence having an extremely high homology. Thus, a natural human FR, obtained for each FR, more than satisfies the consensus sequence of the subgroup though it is derived from different antibodies.

### 3. Natural-sequence humanized antibody

The natural humanized antibody obtained in the present invention comprises human antibody FRs that have been recognized to occur in nature. Though FR1 to FR4 are sometimes derived from different antibodies, homology search between human antibodies permits the selection of the antibodies that only belong to the same subgroup as described above. The FR structure of each antibody in the same subgroup has a structure very similar to

another, and in fact humanized antibodies based on  
consensus sequences in the subgroup have been generated  
(Kettleborough, C. A. et al., Protein Engng. (1991) 4,  
773-783; Satoh, K. et al., Molec. Immun. (1994) 31, 371-  
381).

It is believed that in antibodies, as described  
above, extremely diverse amino acid sequences occur  
naturally through somatic mutation. Only some of the  
structures have been characterized at present. If the FR  
sequence of the antibody obtained cannot be found in  
nature, it is not clear whether the FR is present in  
nature or not. When antibodies are considered as  
pharmaceuticals, the construction of CDR-grafting  
antibody comprising naturally occurring human FRs  
provides such an antibody that has properties superior to  
the conventional humanized antibodies from a viewpoint of  
of the object of the present invention to reduce  
antigenicity.

#### 4. Method of constructing novel humanized antibody

The present invention solves the problem associated  
with humanized antibody constructed by the conventional  
technique of humanization, that is, it eliminates  
antigenicity arising from artificial FRs that are not  
found in nature. Otherwise it is a technology to  
construct humanized antibody by CDR-grafting composed of  
human FRs actually found in nature. The amino acid  
sequences of artificial FRs refer to the amino acid  
sequences of the FRs which as a whole cannot be found in  
nature. The artificial amino acid sequences contained in  
FRs refer to those amino acid sequences that cannot be  
found in nature in FRs.

As the amino acid sequences of FRs that are not  
found in nature, there may be mentioned FRs having an  
amino acid sequence in which human amino acid residues in  
a FR have returned to amino acid residues found in the FR  
of antibody derived from a non-human mammal which is a  
template of humanization in a humanized antibody



constructed by the conventional antibody-humanization technology. Alternatively, in a humanized antibody constructed by the conventional antibody-humanization technology, there may be mentioned FRs having an amino acid sequence that are not found in the antibodies derived from human and non-human mammals.

The method of producing the natural humanized antibody of the present invention is described hereinbelow.

First, a FR of the human antibody for use in CDR-grafting is selected by a conventional technique. The FR is subjected to amino acid substitution to construct a humanized antibody having a biological activity equal to or higher than that of mouse antibody. This is considered as an end product of humanized antibody in the conventional method, but in the present invention it is a mere intermediate for construction of natural humanized antibody having a natural sequence. In the present invention it is called the primary design antibody.

Subsequently, homology search is conducted on each of the FRs of the primary design antibody. FRs having a complete match mean that the FRs have already comprised the natural FRs. On the other hand, a series of natural human FRs are listed that belong to the same subgroup as the primary design antibody and having a homology but not a complete match with the primary design antibody. From the list, there may be selected most appropriate natural human FRs that maintain the amino acid residue of the FR derived from a non-human mammal such as mouse which was important in the construction of the primary design antibody, and that has a homology with the primary design antibody.

Homology search of FRs can be conducted using known databases. Examples of such databases include Swiss Plot, GenBank, PRF, PIR, and GenPept. Homology search is conducted using these databases in which "the FR having a homology with the FR of the primary design antibody"

listed by homology search refers to the FR having a homology in the amino acid sequence of at least 80%, preferably at least 90%, more preferably at least 91%, more preferably at least 92%, more preferably at least 93%, more preferably at least 94%, more preferably at least 95%, more preferably at least 96% or greater, more preferably at least 97% or greater, more preferably at least 98% or greater, and more preferably at least 99% or greater. The homology of protein can be determined by the algorithm described Wilbur, W. J. and Lipman, D. J. Proc. Natl. Acad. Sci. U.S.A. (1983) 80, 726-730.

Amino acid residues of a non-human mammal which were important for construction of the primary design antibody refers to the amino acid residues derived from a non-human FR contained in an artificial FR. Many such amino acid residues are found in the amino acid residues (canonical structure) responsible for the basic structure of antibody molecule, the amino acid residues involved in the maintenance of the structure of CDRs, or the amino acid residues that directly interact with antigen molecule, and include for example an amino acid at position 71 of the H chain, an amino acid at position 94 of the H chain, and the like, though they may vary depending on the antibody.

As mentioned above, if one or a plurality of amino acid residues different between the FR of the primary design antibody and the natural FR are replaced so as to produce humanized antibody having the amino acid residues of a natural human FR, the humanized antibody (natural humanized antibody; termed the secondary design antibody) thus obtained all comprise natural FRs. In this case all human FRs are preferably human FRs that belong to the same subgroup, and more preferably are derived from the same antibody. Furthermore, all human FRs are not required to belong to the same subgroup, as long as they are reshaped into an antibody and provide certain antigen binding activity, and thereby they are not limited to the

human FRs that belong to the same subgroup. According to the present invention, a plurality of amino acid residues mean 2 or more amino acid residues, preferably 2 or more and 10 or less amino acid residues, more preferably 2 or more and 5 or less amino acid residues, more preferably 2 or more and 4 or less amino acid residues, and more preferably 2 or more and 3 or less amino acid residues in the amino acid sequence.

Homology between an artificial FR and a natural human FR is at least 80%, preferably at least 90%, more preferably at least 91%, more preferably at least 92%, more preferably at least 93%, more preferably at least 94%, more preferably at least 95%, more preferably at least 96% or greater, more preferably at least 97% or greater, more preferably at least 98% or greater, and more preferably at least 99% or greater.

Then, the secondary design antibody is allowed to be expressed in a suitable expression system, for example in an animal cell, to evaluate the antigen binding activity, and the like.

Furthermore, the method of preparation of the present invention can be effected even without the actual construction of the primary design antibody. Thus, the primary design antibody is conventionally designed, and without the evaluation thereof the secondary design antibody may be designed, which may be directly evaluated. In fact, however, the identification of important FR residues sometimes involves experiment, and the secondary design antibody is preferably constructed after the conventional primary design antibody has been experimentally constructed.

Specifically, in one aspect of the present invention, the natural humanized antibody of the present invention was produced with mouse anti-HM1.24 antibody (Goto, T. et al., Blood (1994) 84, 1922-1930) as a template.

For natural humanized antibodies designed as

mentioned above, the gene encoding them can be obtained by a known method. For example, several oligonucleotides are synthesized that have overlapping ends corresponding to the DNA encoding the amino acid sequence of the designed natural humanized antibody. A PCR method is carried out using these oligonucleotides as primers. Then, a PCR method is carried out using primers that define the both ends of the DNA encoding the amino acid sequence of the designed natural humanized antibody to obtain the gene encoding the desired natural humanized antibody.

Genes encoding a natural humanized antibody constructed as described above may be expressed in a known method so as to obtain the natural humanized antibody. In the case of mammalian cells, expression may be accomplished using a commonly used useful promoter/enhancer, the antibody gene to be expressed, and DNA in which the poly A signal has been operably linked at 3' downstream thereof, or using a vector containing the same. Examples of the promoter/enhancer include human cytomegalovirus immediate early promoter/enhancer.

Additionally, as the promoter/enhancer which can be used for expression of antibody for use in the present invention, there can be used viral promoters/enhancers such as retrovirus, polyoma virus, adenovirus, and simian virus 40 (SV40), and promoters/enhancers derived from mammalian cells such as human elongation factor 1 $\alpha$  (HEF1 $\alpha$ ).

For example, expression may be readily accomplished by the method of Mulligan et al. (Nature (1979) 277, 108) when the SV40 promoter/enhancer is used, or by the method of Mizushima et al. (Nucleic Acids Res. (1990) 18, 5322) when the HEF1 $\alpha$  promoter/enhancer is used.

In the case of Escherichia coli (E. coli), expression may be effected by operably linking a commonly used useful promoter, a signal sequence for antibody

secretion, and the antibody gene to be expressed, followed by expression thereof. As the promoter, for example, there can be mentioned the lacz promoter and the araB promoter. The method of Ward et al. (Nature (1098) 341, 544-546; FASEB J. (1992) 6, 2422-2427) may be used when lacz promoter is used, and the method of Better et al. (Science (1988) 240, 1041-1043) may be used when araB promoter is used.

As the signal sequence for antibody secretion, when produced in the periplasm of E. coli, the pelB signal sequence (Lei, S.P. et al., J. Bacteriol. (1987) 169, 4379) can be used. After separating the antibody produced in the periplasm, the structure of the antibody is appropriately refolded before use (see, for example, International Patent Publication WO 96/30394, and Japanese Examined Patent Publication (Kokoku) No. 7(1995)-93879).

As the origin of replication, there can be used those derived from SV40, polyoma virus, adenovirus, bovine papilloma virus (BPV) and the like. Furthermore, for the amplification of the gene copy number in the host cell system, expression vectors can include as selectable markers the aminoglycoside transferase (APH) gene, the thymidine kinase (TK) gene, E. coli xanthine guaninephosphoribosyl transferase (Ecogpt) gene, the dihydrofolate reductase (dhfr) gene and the like.

For the production of antibody for use in the present invention, any production system can be used. The production system of antibody preparation comprises the in vitro or the in vivo production system. As the in vitro production system, there can be mentioned a production system which employs eukaryotic cells and the production system which employs prokaryotic cells.

When the eukaryotic cells are used, there are the production systems which employ animal cells, plant cells, and fungal cells. Known animal cells include (1) mammalian cells such as CHO cells (J. Exp. Med. (1995)

108, 945), COS cells, myeloma cells, baby hamster kidney (BHK) cells, HeLa cells, and Vero cells, (2) amphibian cells such as *Xenopus* oocytes (Valle, et al., Nature (1981) 291, 358-340), or (3) insect cells such as sf9, sf21, and Tn5. As CHO cells, preferably dhfr-CHO (Proc. Natl. Acad. Sci. U.S.A. (1980) 77, 4216-4220) that lacks the DHFR gene and CHO K-1 (Proc. Natl. Acad. Sci. U.S.A. (1968) 60, 1275) may be used.

Known plant cells include, for example, those derived from Nicotiana tabacum, which is subjected to callus culture. Known fungal cells include yeasts such as the genus Saccharomyces, for example Saccharomyces cereviceae, or filamentous fungi such as the genus Aspergillus, for example Aspergillus niger.

When the prokaryotic cells are used, there are the production systems which employ bacterial cells. Known bacterial cells include Escherichia coli (E. coli), and Bacillus subtilis.

By transforming these cells with the gene encoding the natural humanized antibody of the present invention and and culturing the transformed cells in vitro, the natural humanized antibody can be obtained. Culturing is carried out in a known method. For example, as the culture liquid, DMEM, MEM, RPMI1640, and IMDM can be used, and serum supplements such as fetal calf serum (FCS) may be used in combination, or serum-free culture medium may be used. In addition, antibodies may be produced in vivo by implanting cells into which the antibody gene has been introduced into the abdominal cavity of an animal and the like.

As in vivo production systems, there can be mentioned those which employ animals and those which employ plants. The gene of antibody is introduced into an animal or a plant, and the antibody is produced in such an animal or a plant and then collected.

When animals are used, there are the production systems which employ mammals and insects.

As mammals, goats, pigs, sheep, mice, and cattle can be used (Vicki Glaser, SPECTRUM Biotechnology Applications, 1993). When mammals are used, transgenic animals can also be used.

5 For example, an antibody gene is inserted into a gene encoding protein which is inherently produced in the milk such as goat  $\beta$  casein to prepare fusion genes. DNA fragments containing the fusion gene into which the antibody gene has been inserted are injected into a goat embryo, and the embryo is introduced into a female goat. 10 The desired antibody is obtained from the milk produced by the transgenic goat borne to the goat who received the embryo or offsprings thereof. In order to increase the amount of milk containing the desired antibody produced 15 by the transgenic goat, hormones may be given to the transgenic goat as appropriate (Ebert, K.M. et al., Bio/Technology (1994) 12, 699-702).

When insects are used, silkworms can be used. When silkworms are used, baculovirus into which the desired 20 antibody gene has been inserted is infected to the silkworm, and the desired antibody can be obtained from the body fluid of the silkworm (Susumu, M. et al., Nature (1985) 315, 592-594).

When plants are used, tobacco, for example, can be 25 used. Moreover, when tobacco is used, the desired antibody gene is inserted into an expression vector for plants, for example pMON 530, and then the vector is introduced into a bacterium such as Agrobacterium tumefaciens. The bacterium is then infected to tobacco 30 such as Nicotiana tabacum to obtain the desired antibody from the leaves of the tobacco (Julian, K.-C. Ma et al., Eur. J. Immunol. (1994) 24, 131-138).

As described above, "hosts" as used herein encompasses animals and plants that produce the desired 35 natural humanized antibody. When antibody is produced in vitro or in vivo production systems, as described above, DNA encoding an H chain or an L chain of an antibody may

002220" 85050550

be separately integrated into an expression vector and a host is transformed simultaneously, or DNA encoding an H chain and DNA encoding an L chain may be integrated into a single expression vector and a host is transformed therewith (see International Patent Publication WO 94-11523).

As method of introducing an expression vector into a host, a known method such as the calcium phosphate method (Virology (1973) 52, 456-467) and the electroporation method (EMBO J. (1982) 1, 841-845) and the like can be used.

A natural humanized antibody produced and expressed as described above can be separated from the inside or outside of the cell or from the host and then may be purified to homogeneity. Separation and purification of the natural humanized antibody for use in the present invention may be accomplished by conventional methods of separation and purification used for protein, without any limitation. Separation and purification may be accomplished by combining, as appropriate, chromatography such as affinity chromatography, filtration, ultrafiltration, salting-out, dialysis and the like (Antibodies: A Laboratory Manual, Ed Harlow and David Lane, Cold Spring Harbor Laboratory, 1988).

As the column used for such affinity chromatography, there can be mentioned Protein A column and Protein G column. As carriers for use in the Protein A column there can be mentioned Hyper D, POROS, Sepharose F.F. (Pharmacia) and the like.

Chromatography other than affinity chromatography includes, for example, ion exchange chromatography, hydrophobic chromatography, gel-filtration, reverse-phase chromatography, adsorption chromatography and the like (Strategies for Protein Purification and Characterization: A Laboratory Course Manual. Ed Daniel R. Marshak et al., Cold Spring Harbor Laboratory Press, 1996).



These chromatographies can be carried out using a liquid chromatography such as HPLC, FPLC, and the like.

5 The concentration of the natural humanized antibody of the present invention can be determined by the measurement of absorbance or by the enzyme-linked immunosorbent assay (ELISA) and the like. Thus, when absorbance measurement is employed, the natural humanized antibody obtained is appropriately diluted with PBS and then the absorbance is measured at 280 nm, followed by  
10 calculation using the absorption coefficient of 1.35 OD at 1 mg/ml.

When the ELISA method is used, measurement is conducted as follows. Thus, 100  $\mu$ l of goat anti-human IgG (manufactured by BIO SOURCE) diluted to 1 mg/ml in 0.1 M  
15 bicarbonate buffer, pH 9.6, is added to a 96-well plate (manufactured by Nunc), and is incubated overnight at 4 °C to immobilize the antibody. After blocking, 100  $\mu$ l each of appropriately diluted natural humanized antibody of the present invention or a sample containing the  
20 antibody, or human IgG (manufactured by CAPPEL) of a known concentration as the standard is added, and incubated at room temperature for 1 hour.

After washing, 100  $\mu$ l of 5000-fold diluted alkaline phosphatase-labeled anti-human IgG antibody (manufactured  
25 by BIO SOURCE) is added, and incubated at room temperature for 1 hour. After washing, the substrate solution is added and incubated, followed by the measurement of absorbance at 405 nm using the MICROPLATE READER Model 3550 (manufactured by Bio-Rad) to calculate  
30 the concentration of the desired antibody. BIAcore (manufactured by Pharmacia) can be used for the measurement of antibody concentration.

The antigen binding activity, binding inhibition activity, and neutralizing activity of the natural  
35 humanized antibody of the present invention can be evaluated by known methods. For example, as methods of

determining the activity of the natural humanized antibody of the present invention, there can be mentioned ELISA, EIA (enzymeimmunoassay), RIA (radioimmunoassay), or fluorescent antibody method. For the evaluation of the above antibody, BIAcore (manufactured by Pharmacia) can be used.

The natural humanized antibody of the present invention may be antibody fragments or modified versions thereof. For example, as fragments of antibody, there may be mentioned Fab, F(ab')<sub>2</sub>, Fv or single-chain Fv (scFv). scFv has a structure in which Fvs of the H chain and the L chain are ligated via a suitable linker.

In order to produce these antibodies, antibodies are treated with an enzyme such as papain or pepsin, or genes encoding these antibody fragments are constructed and then introduced into an expression vector, which is expressed in a suitable host cell to express them (see, for example, Co, M. S. et al., J. Immunol. (1994) 152, 2968-2976; Better, M. and Horwitz, A.H., Methods in Enzymology (1989) 178, 476-496, Academic Press Inc.; Plucktrun, A. and Skerra, A., Methods in Enzymol. (1989) 178, 476-496, Academic Press Inc.; Lamoyi, E., Methods in Enzymol. (1986) 121, 652-663; Rousseaux, J. et al., Methods in Enzymol. (1986) 121, 663-669; Bird, R.E. and Walker, B.W., TIBTECH (1991) 9, 132-137).

scFv can be obtained by ligating the V region of H chain and the V region of L chain of antibody (see, International Patent Publication WO 88-09344). In scFv, the V region of H chain and the V region of L chain are preferably ligated via a linker, preferably a peptide linker (Huston, J.S. et al., Proc. Natl. Acad. Sci. U.S.A. (1988) 85, 5879-5883). The V region of H chain and the V region of L chain in the scFv may be derived from any of the above-mentioned antibodies. As the peptide linker for ligating the V regions, any single-chain peptide comprising, for example, one comprising 12 to 19 amino acid residues may be used (see, United States

Patent No. US 5525491).

5 DNA encoding scFv can be obtained using DNA encoding  
the H chain or the H chain V region of the above antibody  
and DNA encoding the L chain or the L chain V region of  
the above antibody as the template by amplifying the  
portion of the DNA encoding the desired amino acid  
sequence among the above sequences by the PCR technique  
with the primer pair specifying the both ends thereof,  
and by further amplifying the combination of DNA encoding  
10 the peptide linker portion and the primer pair which  
defines that both ends of said DNA be ligated to the H  
chain and the L chain, respectively.

Once DNAs encoding scFv are constructed, an  
expression vector containing them and a host transformed  
15 with said expression vector can be obtained by the  
conventional methods, and scFv can be obtained using the  
resultant host by the conventional methods.

These antibody fragments can be produced by  
obtaining the gene thereof in a similar manner to that  
20 mentioned above and by allowing it to be expressed in a  
host. "Antibody" as used in the claim of the present  
application encompasses these antibody fragments.

As modified antibodies, antibodies associated with  
various molecules such as polyethylene glycol (PEG) can  
25 be used. "Antibody" as used in the claim of the present  
application encompasses these modified antibodies. These  
modified antibodies can be obtained by chemically  
modifying the antibodies thus obtained. These methods  
have already been established in the art.

30 The natural humanized antibody of the present  
invention may be administered orally or parenterally,  
either systemically or topically. The parenteral route  
may be selected from intravenous injection such as drip  
infusion, intramuscular injection, intraperitoneal  
35 injection, and subcutaneous injection, and the method of  
administration may be chosen, as appropriate, depending  
on the age and the condition of the patient.

5 The natural humanized antibody of the present invention may be administered at a dosage that is sufficient to treat or to block at least partially the pathological condition. For example, the effective dosage is chosen from the range of 0.01 mg to 100 mg per kg of body weight per administration. Alternatively, the dosage in the range of 1 to 1000 mg, preferably 5 to 50 mg per patient may be chosen. However, the natural humanized antibody of the present invention is not limited to these dosages.

10 The natural humanized antibody of the present invention may contain pharmaceutically acceptable carriers or additives depending on the route of administration. Examples of such carriers or additives include water, a pharmaceutical acceptable organic solvent, collagen, polyvinyl alcohol, polyvinylpyrrolidone, a carboxyvinyl polymer, carboxymethyl cellulose sodium, polyacrylic sodium, sodium alginate, water-soluble dextran, carboxymethyl starch sodium, pectin, methyl cellulose, ethyl cellulose, xanthan gum, gum Arabic, casein, gelatin, agar, diglycerin, propylene glycol, polyethylene glycol, Vaseline, paraffin, stearyl alcohol, searic acid, human serum albumin (HSA), mannitol, sorbitol, lactose, a pharmaceutically acceptable surfactant and the like. Additives used are chosen from, but not limited to, the above or combinations thereof depending on the dosage form.

### 30 Reference Examples

Before explaining the present invention with reference to the working examples, reference examples will be described as the premise thereof.

#### 35 Reference example 1. Cloning of cDNA encoding the variable region of a mouse anti-HM1.24 antibody

##### 1. Isolation of messenger RNA (mRNA)

Using the Fast Track mRNA Isolation Kit Version

3.2 (manufactured by Invitrogen) according to the instruction attached thereto, mRNA was isolated from  $2 \times 10^8$  hybridoma cells (FERM BP-5233) that produce a mouse anti-HM1.24 antibody.

5           2. Amplification of the gene encoding the variable region of antibody by the PCR method

        PCR was carried out using the amplification Thermal Cycler (manufactured by Perkin Elmer Cetus).

10           2-1. Amplification and fragmentation of the gene encoding the V region of a mouse L chain

        From the mRNA thus isolated, single stranded cDNA was synthesized using the AMV Reverse Transcriptase First-strand cDNA Synthesis Kit (manufactured by Life Science) and used for PCR. As primers used for PCR, MKV (Mouse Kappa Variable) primers (Jones, S.T. et al, 15 Bio/Technology, 9, 88-89, (1991)) shown in SEQ ID NO: 29 to 39 that hybridize with the leader sequence of a mouse kappa type L chain were used.

        A hundred microliters of the PCR solution 20 containing 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 0.1 mM dNTPs (dATP, dGTP, dCTP, dTTP), 1.5 mM MgCl<sub>2</sub>, 5 units of DNA polymerase Ampli Taq (manufactured by Perkin Elmer Cetus), 0.25 mM of the MKV primers shown in SEQ ID NO: 29 to 39, 3 mM of the MKC primer shown in SEQ ID NO: 40, and 25 100 ng of single stranded cDNA was covered with 50  $\mu$ l of a mineral oil, and then heated at an initial temperature of 94 °C for 3 minutes, and then at 94 °C for 1 minute, at 55 °C for 1 minute, and at 72 °C for 1 minute in this order. After repeating this cycle for 30 times, the 30 reaction mixture was incubated at 72 °C for 10 minutes. The amplified DNA fragment was purified by the low melting point agarose (manufactured by Sigma), and digested with XmaI (manufactured by New England Biolabs) and SalI (manufactured by Takara Shuzo) at 37°C.

35           2-2. Amplification and fragmentation of cDNA encoding the V region of a mouse H chain

        The gene encoding the V region of a mouse H

chain was amplified by the 5'-RACE method (Rapid Amplification of cDNA ends; Frohman, M.A. et al., Proc. Natl. Acad. Sci. USA, 85, 8998-9002, (1988), Edwards, J.B.D.M., et al., Nucleic Acids Res., 19, 5227-5232, (1991)). After cDNA was synthesized using primer P1 (SEQ ID NO: 63) that specifically hybridizes with the constant region of mouse IgG2a, cDNA encoding the V region of a mouse H chain was amplified by the 5'-AmplifINDER RACE KIT (manufactured by CLONTECH) using the primer MHC 2a (SEQ ID NO: 64) that specifically hybridizes with the constant region of mouse IgG2a and the anchor primer (SEQ ID NO: 101) attached to the kit. The amplified DNA fragment was purified with the low melting point agarose (manufactured by Sigma) and digested with EcoRI (manufactured by Takara) and XmaI (manufactured by New England Biolabs) at 37°C.

### 3. Linking and transformation

The DNA fragment comprising the gene encoding the V region of the mouse kappa type L chain prepared as above was ligated to the pUC19 vector prepared by digesting with SalI and XmaI by reacting in a reaction mixture containing 50 mM Tris-HCl (pH 7.6), 10 mM MgCl<sub>2</sub>, 10 mM dithiothreitol, 1 mM ATP, 50 mg/ml of polyethylene glycol (8000) and one unit of T4 DNA ligase (manufactured by GIBCO-BRL) at 16 °C for 2.5 hours. Similarly, the gene encoding the V region of the mouse H chain was reacted and ligated to pUC19 vector prepared by digesting with EcoRI and XmaI at 16 °C for three hours.

Then 10 µl of the above ligation mixture was added to 50 µl of the competent cells of Escherichia coli DH5 , which was left on ice for 30 minutes, at 42 °C for one minute, and again on ice for one minute.

Subsequently 400 µl of 2xYT medium (Molecular Cloning: A Laboratory Manual, Sambrook et al., Cold Spring Harbor Laboratory Press, (1989)) was added thereto, incubated at 37°C for one hour, and then the E. coli was plated on the

2xYT agar medium (Molecular Cloning: A Laboratory Manual, Sambrook et al., Cold Spring Harbor Laboratory Press, (1989)) containing 50 µg/ml of ampicillin, and then incubated overnight at 37°C to obtain the E. coli transformant.

The transformant was cultured overnight at 37°C in 10 ml of the 2xYT medium containing 50 µg/ml of ampicillin, and then from this culture plasmid DNA was prepared using the alkali method (Molecular Cloning: A Laboratory Manual, Sambrook et al., Cold Spring Harbor Laboratory Press, (1989)).

The plasmid thus obtained containing the gene encoding the V region of the mouse kappa type L chain derived from the hybridoma that produces the anti-HM1.24 antibody was termed pUCHMVL9. The plasmid obtained in the above-mentioned method containing the gene encoding the V region of the mouse H chain derived from the hybridoma that produces the anti-HM1.24 antibody was termed pUCHMVHR16.

Reference Example 2. Determination of the nucleotide sequence of DNA

The nucleotide sequence of the cDNA coding region in the above-mentioned plasmid was determined using the automatic DNA sequencer (manufactured by Applied Biosystem Inc.) and Taq Dye Deoxy Terminator Cycle Sequencing Kit (manufactured by Applied Biosystem Inc.) in the protocol indicated by the manufacturer.

The nucleotide sequence of the gene encoding the V region of the L chain of the mouse anti-HM1.24 antibody contained in the plasmid pUCHMVL9 is shown in SEQ ID NO: 1. The nucleotide sequence of the gene encoding the V region of the H chain of the mouse anti-HM1.24 antibody contained in the plasmid pUCHMVHR16 is shown in SEQ ID NO: 3.

Reference Example 3. Determination of CDR

The overall structures of the V regions of an L

chain and an H chain have a similarity with each other in which four framework portions are linked by three hypervariable regions, i.e. complementarity determining regions (CDR). The amino acid sequence of the framework is relatively well conserved but variation in the amino acid sequence is extremely high (Kabat, E.A., et al., "Sequences of Proteins of Immunological Interest", US Dept. Health and Human Services, 1983).

Based on these facts, the amino acid sequence of the variable region of the anti-HM1.24 antibody was fitted to the database of the amino acid sequences of antibodies to investigate homology, and the CDR region was determined as shown in Table 1.

Table 1

Plasmid	Sequence No.	CDR(1)	CDR(2)	CDR(3)
pUCHMVL9	5 to 7	24-34	50-56	89-97
pUCHMVHR16	8 to 10	31-35	50-66	99-109

Reference Example 4. Confirmation of expression of the cloned cDNA (Construction of the chimera anti-HM1.24 antibody)

1. Construction of an expression vector

In order to construct an expression vector that expresses a chimera anti-HM1.24 antibody, cDNA clones pUCHMVL9 and pUCHMVHR16 encoding the V regions of the L chain and the H chain of the mouse anti-HM1.24 antibody, respectively, were modified by the PCR method, and then introduced into the HEF expression vector (International Patent Publication No. WO 92-19759).

The backward primer ONS-L722S (SEQ ID NO: 65) for the V region of an L chain and the backward primer VHR16S (SEQ ID NO: 66) for the V region of an H chain were designed so that they hybridize to the DNA encoding the start of the leader sequence of the V region of each and they have the Kozak consensus sequence (Kozak, M. et al., J. Mol. Biol., 196, 947-950, (1987)) and the recognition site for HindIII restriction enzyme. The forward primer VL9A (SEQ ID NO: 67) for the V region of



an L chain and the forward primer VHR16A (SEQ ID NO: 68) for the V region of an H chain were designed so that they hybridize to the DNA sequence encoding the end of the J region and they have a splice donor sequence and the recognition site for BamHI restriction enzyme.

One hundred µl of the PCR reaction mixture containing 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 0.1 mM dNTPs , 1.5 mM MgCl<sub>2</sub>, 100 pmole each of each primer, 100 ng of template DNA (pUCHMVL9 or pUCHMVHR16), and 5 units of Ampli Taq enzyme was covered with 50 µl of a mineral oil, and then after the initial denaturation at 94 °C, heated at 94 °C for 1 minute, at 55 °C for 1 minute and at 72 °C for 1 minute for 30 cycles and finally incubated at 72 °C for 10 minutes.

The PCR product was purified by the low melting point agarose gel, and digested with HindIII and BamHI, and then cloned to HEF-VL-gk for the V region of the L chain and to HEF-VH-gyl for the V region of the H chain. After determination of the DNA sequence, the plasmids containing the DNA fragment that contains the correct DNA sequence were designated as HEF-1.24L-gk and HEF-1.24H-gyl, respectively.

The regions encoding the respective variable region from the above plasmids HEF-1.24L-gk and HEF-1.24H-gyl were digested with restriction enzymes HindIII and BamHI to make restriction fragments, which were inserted to the HindIII site and the BamHI sites of plasmid vector pUC19 and they were designated as pUC19-1.24L-gk and pUC19-1.24H-gyl, respectively.

Escherichia coli containing respective plasmids pUC19-1.24L-gk and pUC19-1.24H-gyl were designated as Escherichia coli DH5 (pUC19-1.24L-gk) and Escherichia coli DH5 (pUC19-1.24H-gyl), and were internationally deposited on August 29, 1996, with the National Institute

of Bioscience and Human-Technology, Agency of Industrial  
Science and Technology, MITI (Higashi 1-Chome 1-3,  
Tsukuba city, Ibalaki prefecture, Japan) under the  
accession numbers FERM BP-5646 and FERM BP-5644,  
5 respectively, under the provisions of the Budapest  
Treaty.

## 2. Transfection into COS-7 cells

In order to observe the transient expression  
of the chimera anti-HM1.24 antibody, the above expression  
10 vectors were tested in the COS-7 (ATCC CRL-1651) cells.  
HEF-1.24L-gk and HEF-1.24H-gyl were cotransformed into  
COS-7 cells by electroporation using the Gene Pulser  
instrument (manufactured by BioRad). Each DNA (10 µg)  
was added to 0.8 ml aliquots of  $1 \times 10^7$  cells/ml in PBS,  
15 and was subjected to pulses at 1500 V and a capacity of  
25 µF.

After a recovery period of 10 minutes at room  
temperature, the electroporated cells were added to 30 ml  
of the DHEM culture liquid (manufactured by GIBCO)  
20 containing 10% γ-globulin-free bovine fetal serum. After  
incubation of 72 hours in the CO<sub>2</sub> incubator BNA120D  
(manufactured by TABAI), the culture supernatant was  
collected, the cell debris was removed by centrifugation,  
and the supernatant was used for the following  
25 experiment.

## 3. FCM analysis

The antigen binding activity of the chimera  
anti-HM1.24 antibody was investigated by FCM (flow  
cytometry) analysis using the KPMM2 cells. After  $4.7 \times$   
30  $10^5$  KPMM2 cells (Japanese Unexamined Patent Publication  
(Kokai) No. 7(1995)-236475) were washed with PBS(-), 50  
µl of the culture of COS-7 cells that produce the  
above-mentioned chimera anti-HM1.24 antibody and 50 µl of  
FACS buffer (PBS(-) containing 2% bovine fetal serum and  
35 0.1% sodium azide), or 5 µl of 500 µg/ml purified mouse

anti-HM1.24 antibody and 95 µl of the FACS buffer were added, and incubated at the temperature of ice for one hour.

As a control, 50 µl of 2 µg/ml chimera SK2 (International Patent Publication No. WO 94-28159) and 50 µl of the FACS buffer, or 5 µl of 500 µg/ml purified mouse IgG2ak (UPC10) (manufactured by CAPPEL ) instead of purified mouse anti-HM1.24 antibody, and 95 µl of FACS buffer were added, and similarly incubated. After washing with the FACS buffer, 100 µl of 25 µg/ml FITC-labeled goat anti-human antibody (GAH) (manufactured by CAPPEL) or 10 µg/ml FITC labeled goat anti-mouse antibody (GAM) (manufactured by Becton Dickinson) were added, and incubated at a temperature of ice for 30 minutes. After washing with the FACS buffer, it was suspended in one ml of the FACS buffer, and fluorescence intensity of each cell was measured by the FACScan (manufactured by Becton Dickinson).

As shown in Fig. 1, it was revealed that the chimera anti-HM1.24 antibody bound to the KPMM2 cell because the peak of fluorescence intensity shifted to the right in the chimera anti-HM1.24 antibody-added cells as compared to the control similarly to the case where mouse anti-HM1.24 antibody was added. This confirmed that the cloned cDNA encodes the variable region of the mouse anti-HM1.24 antibody.

Reference Example 5. Establishment of the CHO cell line that stably produces a chimera anti-HM1.24 antibody

1. Construction of an expression vector for the chimera H chain

By digesting the above plasmid HEF-1.24H-gyl with the restriction enzymes PvuI and BamHI, an about 2.8 kbp fragment containing the EF1 promoter and the DNA encoding the V region of the H chain of the mouse

anti-HM1.24 antibody was purified using 1.5% low melting point agarose gel. Then, the above DNA fragment was inserted into an about 6 kbp fragment prepared by digesting the expression vector used for a human H chain expression vector, DHFR- $\Delta$ E-Rvh-PM1f (see International Patent Publication No. WO 92/19759), containing the DHFR gene and the gene encoding the constant region of a human H chain with PvuI and BamHI to construct an expression vector, DHFR- $\Delta$ E-HEF-1.24-H-gyl, for the H chain of the chimera anti-HM1.24 antibody.

## 2. Gene introduction into CHO cells

In order to establish a stable production system of the chimera anti-HM1.24 antibody, the genes of the above-mentioned expression vectors, HEF-1.24L-gk and DHFR- $\Delta$ E-HEF-1.24H-gyl, that were linearized by digestion with PvuI were simultaneously introduced into the CHO cell DXBII (donated from the Medical Research Council Collaboration Center) by the electroporation method under the condition similar to the above-mentioned one (the above-mentioned transfection into the COS-7 cells).

## 3. Gene amplification by MTX

Among the gene-introduced CHO cells, only those CHO cells in which both of the L chain and the H chain expression vectors have been introduced can survive in the nucleoside-free  $\alpha$ -MEM culture liquid (manufactured by GIBCO-BRL) to which 500  $\mu$ g/ml G418 (manufactured by GIBCO-BRL) and 10% bovine fetal serum were added, and so they were selected. Subsequently, 10 nM MTX (manufactured by Sigma) was added to the above culture liquid. Among the clones that propagated, those that produce the chimera anti-HM1.24 antibody in large amounts were selected. As a result, clones #8 to #13 that exhibited a production efficiency of about 20  $\mu$ g/ml of the chimera antibody were obtained and termed the chimera anti-HM1.24 antibody-producing cell lines.

Reference Example 6. Construction of the chimera  
anti-HM1.24 antibody

5 The chimera anti-HM1.24 antibody was  
constructed in the following method. The above chimera  
anti-HM1.24 antibody-producing CHO cells were subjected  
to continuous culture for 30 days using as the medium  
Iscove's Modified Dulbecco's Medium (manufactured by  
GIBCO-BRL) containing 5%  $\gamma$ -globulin-free newborn bovine  
serum (manufactured by GIBCO-BRL) by the high-density  
10 cell culture instrument Verax system 20 (manufactured by  
CELLEX BIOSCIENCE Inc.).

On day 13, 20, 23, 26, and 30 after starting  
the culture, the culture liquid was recovered using a  
pressurized filter unit SARTOBRAN (manufactured by  
15 Sartorius), and then the chimera anti-HM1.24 antibody was  
affinity-purified using a large-volume antibody  
collection system Afi-Prep System (manufactured by Nippon  
Gaishi) and Super Protein A column (bed volume: 100 ml,  
manufactured by Nippon Gaishi) using PBS as the  
20 absorption/wash buffer and 0.1 M sodium citrate buffer  
(pH 3) as the elution buffer according to the attached  
instructions. The eluted fractions were adjusted to  
about pH 7.4 by immediately adding 1 M Tris-HCl (pH 8.0).  
Antibody concentration was measured by absorbance at 280  
25 nm and calculated with 1  $\mu$ g/ml as 1.35 OD.

Reference Example 7. Determination of activity of  
the chimera anti-HM1.24 antibody

Chimera anti-HM1.24 antibody was evaluated by  
the following binding inhibition activity.

- 30 1. Measurement of binding inhibition activity  
1-1. Construction of a biotinylated anti-HM1.24  
antibody

After the mouse anti-HM1.24 antibody was  
diluted with 0.1 M bicarbonate buffer to 4 mg/ml, 4  $\mu$ l of  
35 50 mg/ml Biotin-N-hydroxy succinimide (manufactured by EY  
LABS Inc.) was added and reacted at room temperature for

3 hours. Thereafter, 1.5 ml of 0.2 M glycine solution was added thereto, incubated at room temperature for 30 minutes to stop the reaction, and then the biotinylated IgG fractions were collected using the PD-10 column (manufactured by Pharmacia Biotech).

1-2. Measurement of binding inhibition activity

The binding inhibition activity by the biotin-labeled mouse anti-HM1.24 antibody was measured by the Cell-ELISA using the human amniotic membrane cell line WISH cells (ATCC CCL 25). The Cell-ELISA plates were prepared as follows. To a 96-well plate was added  $4 \times 10^5$  cells/ml prepared with PRMI 1640 medium supplemented with 10% fetal bovine serum, incubated overnight, and after washing twice with PBS(-), were immobilized with 0.1% glutaraldehyde (manufactured by Nacalai Tesque Inc.).

After blocking, 50  $\mu$ l of serial dilutions of the chimera anti-HM1.24 antibody or the mouse anti-HM1.24 antibody obtained by affinity purification was added to each well and simultaneously 50  $\mu$ l of 2  $\mu$ g/ml biotin-labeled mouse anti-HM1.24 antibody was added, incubated at room temperature for two hours, and then the peroxidase-labeled streptavidin (manufactured by DAKO) was added. After incubating at room temperature for one hour and then washing, the substrate solution was added. After stopping the reaction by adding 50  $\mu$ l of 6N sulfuric acid, absorbance at 490 nm was measured using the MICROPLATE READER Model 3550 (manufactured by Bio-Rad).

The result, as shown in Fig. 2, revealed that the chimera anti-HM1.24 antibody has a similar binding inhibition activity with the mouse anti-HM1.24 antibody as the biotin-labeled mouse anti-HM1.24 antibody. This indicates that the chimera antibody had the same V region as the mouse anti-HM1.24 antibody.

Reference Example 8. Measurement of the ADCC  
activity of the chimera anti-HM1.24 antibody

ADCC (Antibody-dependent Cellular Cytotoxicity) activity was measured according to the method as set forth in Current Protocols in Immunology, Chapter 7, Immunologic studies in humans, Editor, Johan E. Coligan et al., John Wiley & Sons, Inc., 1993.

1. Preparation of effector cells

Monocytes were separated from the peripheral blood or bone marrow of healthy humans and patients with multiple myeloma by the density centrifugation method. Thus, an equal amount of PBS(-) was added to the peripheral blood and the bone marrow of healthy humans and patients with multiple myeloma, which was layered on Ficoll (manufactured by Pharmacia)-Conrey (manufactured by Daiichi Pharmaceutical Co. Ltd.) (specific gravity, 1.077), and was centrifuged at 400 g for 30 minutes. The monocyte layer was collected, and washed twice with RPMI 1640 (manufactured by Sigma) supplemented with 10% bovine fetal serum (manufactured by Witaker), and prepared at a cell density of  $5 \times 10^6$ /ml with the same culture liquid.

2. Preparation of target cells

The human myeloma cell line RPMI 8226 (ATCC CCL 155) was radiolabeled by incubating in the RPMI 1640 (manufactured by Sigma) supplemented with 10% bovine fetal serum (manufactured by Witaker) together with 0.1 mCi of  $^{51}\text{Cr}$ -sodium chromate at 37 °C for 60 minutes. After radiolabeling, cells were washed three times with Hanks balanced salt solution (HBSS) and adjusted to a concentration of  $2 \times 10^5$ /ml.

3. ADCC assay

Into a 96-well U-bottomed plate (manufactured by Corning) were added 50  $\mu\text{l}$  of  $2 \times 10^5$  target cells/ml, 1  $\mu\text{g}/\text{ml}$  of affinity-purified chimera anti-HM1.24 antibody and mouse anti-HM1.24 antibody, or control human IgG (manufactured by Serotec), and the plate was held at

4 °C for 15 minutes.

Then, 100 µl of  $5 \times 10^5$  effector cells/ml was added thereto, and the result was cultured in a CO<sub>2</sub> incubator for 4 hours, whereupon the ratio (E:T) of the effector cells (E) to the target cells (T) was set at 0:1, 5:1, 20:1, or 50:1.

One hundred µl of the supernatant was taken and the radioactivity released into the culture supernatant was measured by a gamma counter (ARC361, manufactured by Aloka). For measurement of the maximum radioactivity, 1% NP-40 (manufactured by BRL) was used. Cytotoxicity (%) was calculated by  $(A-C)/(B-C) \times 100$ , wherein A is radioactivity (cpm) released in the presence of antibody, B is radioactivity (cpm) released by NP-40, and C is radioactivity (cpm) released by the culture liquid alone without antibody.

As shown in Fig. 3, when the chimera anti-HM1.24 antibody was added as compared to the control IgG1, cytotoxicity increased with the increase in the E:T ratio, which indicated that this chimera anti-HM1.24 antibody has ADCC activity. Furthermore, since there was no cytotoxicity observed even when the mouse anti-HM1.24 antibody was added, it was shown that the Fc portion of human antibody is required to obtain ADCC activity when the effector cell is a human-derived cell.

Reference Example 9. Construction of the reshaped human anti-HM1.24 antibody

1. Designing of the V region of the reshaped human anti-HM1.24 antibody

In order to construct the reshaped human antibody in which the CDR of mouse monoclonal antibody has been transplanted to a human antibody, it is preferred that there is a high homology between the FR of the mouse antibody and the FR of the human antibody. Thus, the V regions of the L chain and the H chain of the mouse anti-HM1.24 antibody were compared to the V regions



of all known antibodies whose structure has been elucidated using the Protein Data Bank.

5 The V region of the L chain of the mouse anti-HM1.24 antibody is most similar to the consensus sequence of the subgroup IV (HSGIV) of the V region of a human L chain with a homology of 66.4%. On the other hand, it has shown a homology of 56.9%, 55.8%, and 61.5% with HSGI, HSGII and HSG III, respectively.

10 When the V region of the L chain of the mouse anti-HM1.24 antibody is compared to the V region of the L chain of known human antibodies, it has shown a homology of 67.0% with the V region REI of a human L chain, one of the subgroups I of the V region of a human L chain. Thus, the FR of REI was used as the starting material for  
15 construction of the V region of the L chain of the reshaped human anti-HM1.24 antibody.

Version a of the L chain V region of the reshaped human anti-HM1.24 antibody was designed. In this version, human FR was made identical with the  
20 REI-based FR present in the reshaped human CAMPATH-1H antibody (see Riechmann, L. et al., Nature 322, 21-25, (1988), the FR contained in version a of the V region of the L chain of the reshaped human anti PM-1 antibody described in International Patent Publication No. WO  
25 92-19759), and the mouse CDR was made identical with the CDR in the V region of the L chain of the mouse anti-HM1.24 antibody.

The H chain V region of the mouse anti-HM1.24 antibody is most similar to the consensus sequence of  
30 HSGI of the V region of a human H chain with a homology of 54.7%. On the other hand, it shows a homology of 34.6% and 48.1% with HSGII and HSGIII, respectively. When the V region of the H chain of the mouse anti-HM1.24 antibody is compared to the V region of the H chain of  
35 known human antibodies, FR1 to FR3 were most similar to the V region of the H chain of the human antibody HG3, one of subgroup I of the V region of a human H chain

(Rechavi, G. et al., Proc. Natl. Acad. Sci. USA, 80, 855-859), with a homology of 67.3%.

Therefore, the FR of the human antibody HG3 was used as the starting material for construction of the V region of the H chain of the reshaped human anti-HM1.24 antibody. However, since the amino acid sequence of the FR4 of human HG3 has not been described, the amino acid sequence of the FR4 of the human antibody JH6 (Ravetch, J.V. et al., Cell, 27, 583-591) that shows the highest homology with the FR4 of the H chain of the mouse anti-HM1.24 antibody was used. The FR4 of JH6 has the same amino acid sequence as that of the FR4 of the H chain of the mouse anti-HM1.24 antibody except for one amino acid.

In the first version a of the V region of the H chain of the reshaped human anti-HM1.24 antibody, FR1 to FR3 were made identical with the FR1 to FR3 of human HG3, and the CDR was made identical with the CDR of the V region of the H chain of the mouse anti-HM1.24 antibody, except that the amino acids at position 30 in the human FR1 and position 71 in the human FR3 were made identical with the amino acids in the mouse anti-HM1.24 antibody.

## 2. Construction of the V region of the L chain of the reshaped human anti-HM1.24 antibody

The L chain of the reshaped human anti-HM1.24 antibody was constructed by the CDR grafting in the PCR method. The method is shown in Fig. 4. Eight PCR primers were used for construction of the reshaped human anti-HM1.24 antibody (version a) having the FR derived from the human antibody REI. The external primers A (SEQ ID NO: 69) and H (SEQ ID NO: 70) were designed to hybridize with the DNA sequence of the expression vector HEF-VL-gk.

The CDR grafting primers L1S (SEQ ID NO: 71), L2S (SEQ ID NO: 72), and L3S (SEQ ID NO: 73) have the sense DNA sequence. The CDR grafting primers L1A (SEQ ID NO: 74), L2A (SEQ ID NO: 75), and L3A (SEQ ID NO: 76)

have the antisense DNA sequence, each having a complementary DNA sequence (20 to 23 bp) to the DNA sequence at the 5'-end of the primers L1S, L2S, and L3S, respectively.

5           In the first stage of PCR, the four reactions A-L1A, L1S-L2A, L2S-L3A, and L3S-H were conducted to purify each PCR product. The four PCR products from the first PCR were allowed to assemble with one another by their own complementarity (see International Patent  
10       Publication No. WO 92-19759). Then, external primers A and H were added to amplify the full-length DNA encoding the V region of the L chain of the reshaped human anti-HM1.24 antibody (the second PCR). In the  
15       above-mentioned PCR, the plasmid HEF-RVL-M21a (see International Patent Publication No. WO 95-14041) encoding the version a of the V region of the L chain of the reshaped human ONS-M21 antibody based on the human antibody REI-derived FR was employed as a template.

20           In the first stage of PCR, the PCR mixture containing 10 mM Tris-HCl (pH 8.3), 50 mM KCl, 0.1 mM dNTPs, 1.5 mM MgCl<sub>2</sub>, 100 ng of template DNA, 100 pmole of each primer, and 5 u of Ampli Taq was used. Each PCR tube was covered with 50 µl of a mineral oil. Then after  
25       it was first denatured by heating at 94 °C, it was subjected to a reaction cycle of 94 °C for 1 minute, 55 °C for 1 minute and 72 °C for 1 minute, and then was incubated at 72 °C for 10 minutes.

30           PCR products A-L1A (215 bp), L1S-L2A(98 bp), L2S-L3A (140 bp), and L3S-H (151 bp) were purified using 1.5% low melting point agarose gel and were assembled in the second PCR. In the second PCR, 98 µl of PCR mixture containing 1 µg each of the first stage PCR products and  
35       5 u of Ample Taq was incubated for 2 cycles of 94 °C for 2 minutes, 55 °C for 2 minutes, and 72 °C for 2 minutes, and then 100 pmole each of the external primers (A and H) was added. The PCR tube was coated with 50 µl of a

mineral oil and 30 cycles of PCR were conducted under the same condition as above.

5 A 516 bp DNA fragment resulting from the second PCR was purified using 1.5% low melting point agarose gel, digested with BamHI and HindIII, and the DNA fragments thus obtained were cloned into the HEF expression vector HEF-VL-gk. After determining the DNA sequence, the DNA fragment having the correct amino acid sequence of the V region of the L chain of the reshaped  
10 human anti-HM1.24 antibody was designated as plasmid HEF-RVLa-AHM-gk. The amino acid sequence and the nucleotide sequence of the V region of L chain contained in this plasmid HEF-RVLa-AHM-gk are shown in SEQ ID NO: 11.

15 The version b of the V region of the L chain of the reshaped human anti-HM1.24 antibody was constructed by mutagenesis using PCR. Mutagen primers FTY-1 (SEQ ID NO: 77) and FTY-2 (SEQ ID NO: 78) were so designed as to mutate phenylalanine at position 71 to tyrosine.

20 After the above primers were amplified using the plasmid HEF-RVLa-AHM-gk as a template, the final product was purified by digesting with BamHI and HindIII. The DNA fragments obtained were cloned into the HEF expression vector HEF-VL-gk to obtain plasmid  
25 HEF-RVLb-AHM-gk. The amino acid sequence and the base sequence of the V region of the L chain contained in this plasmid HEF-RVLb-AHM-gk are shown in SEQ ID NO: 13.

3. Construction of the H chain V region of the reshaped human anti-HM1.24 antibody

30 3-1. Construction of versions a to e of the H chain V region of the reshaped human anti-HM1.24 antibody

DNA encoding the V region of the H chain of the reshaped human anti-HM1.24 antibody was designed as follows. By linking the DNA sequence encoding the FR1 to  
35 3 of the human antibody HG3 and the FR4 of the human

antibody JH6 to the DNA sequence encoding the CDR of the V region of the H chain of the mouse anti-HM1.24 antibody, the full length DNA encoding the V region of the H chain of the reshaped human anti-HM1.24 antibody was designed.

Then, to the 5'-end and the 3'-end of this DNA sequence the HindIII recognition site/KOZAK consensus sequence and BamHI recognition site/splice donor sequence, respectively, were attached so as to enable insertion of the HEF expression vector.

The DNA sequence thus designed was divided into four oligonucleotides. Subsequently, oligonucleotides which potentially hinder assembly of these oligonucleotides were subjected to computer analysis for the secondary structure. The sequences of the four oligonucleotides RVH1 to RVH4 are shown in SEQ ID NO: 79 to 82. These oligonucleotides have a length of 119 to 144 bases and have the 25 to 26 bp overlapping region. Among the oligonucleotides, RVH2 (SEQ ID NO: 80) and RVH4 (SEQ ID NO: 82) have the sense DNA sequence, and RVH1 (SEQ ID NO: 79) and RVH3 (SEQ ID NO: 81) have the antisense DNA sequence. The method for assembling these four oligonucleotides by the PCR method is shown in the figure (see Fig. 5).

The PCR mixture (98  $\mu$ l) containing 100 ng each of the four oligonucleotides and 5 u of Ampli Taq was first denatured by heating at 94 °C for 2 minutes, and was subjected to two cycles of incubation comprising 94 °C for 2 minutes, 55 °C for 2 minutes and 72 °C for 2 minutes. After 100 pmole each of RHP1 (SEQ ID NO: 83) and RHP2 (SEQ ID NO: 84) were added as the external primer, the PCR tube was coated with 50  $\mu$ l of a mineral oil. Then it was first denatured by heating at 94 °C for 1 minute, and then was subjected to 38 cycles of 94 °C for 1 minute, 55 °C for 1 minute and 72 °C for 1 minute, and then was incubated at 72 °C for 10 minutes.

The 438 bp DNA fragment was purified using 1.5% low melting point agarose gel, digested with HindIII and BamHI, and then cloned into the HEF expression vector HEF-VH-gy1. After determination of the base sequence, the plasmid that contains the DNA fragment encoding the amino acid sequence of the correct V region of the H chain was designated as HEF-RVHa-AHM-gy1. The amino acid sequence and the base sequence of the V region of the H chain contained in this plasmid HEF-RVHa-AHM-gy1 are shown in SEQ ID NO: 11.

Each of versions b, c, d, and e of the V region of the H chain of the reshaped human anti-HM1.24 antibody was constructed as follows.

Using as the mutagen primer BS (SEQ ID NO: 85) and BA (SEQ ID NO: 86) designed to mutate arginine at position 66 to lysine and, as a template DNA, the plasmid HEF-RVHa-AHM-gy1 by the PCR method, version b was amplified to obtain plasmid HEF-RVHb-AHM-gy1. The amino acid sequence and the base sequence of the V region of the H chain contained in this plasmid HEF-RVHb-AHM-gy1 are shown in SEQ ID NO: 17.

Using as the mutagen primer CS (SEQ ID NO: 87) and CA (SEQ ID NO: 88) designed to mutate threonine at position 73 to lysine and, as a template DNA, the plasmid HEF-RVHa-AHM-gy1 by the PCR method, version c was amplified to obtain plasmid HEF-RVHc-AHM-gy1. The amino acid sequence and the base sequence of the V region of the H chain contained in this plasmid HEF-RVHc-AHM-gy1 are shown in SEQ ID NO: 19.

Using as the mutagen primer DS (SEQ ID NO: 89) and DA (SEQ ID NO: 90) designed to mutate arginine at position 66 to lysine and threonine at position 73 to lysine and as a template DNA the plasmid HEF-RVHa-AHM-gy1 by the PCR method, version d was amplified to obtain

plasmid HEF-RVHd-AHM-gyl. The amino acid sequence and the base sequence of the V region of the H chain contained in this plasmid HEF-RVHd-AHM-gyl are shown in SEQ ID NO: 21.

5                   Using as the mutagen primer ES (SEQ ID NO: 91) and EA (SEQ ID NO: 92) designed to mutate valine at position 67 to alanine and methionine at position 69 to leucine and as a template DNA the plasmid HEF-RVHa-AHM-gyl, version e was amplified to obtain

10                   plasmid HEF-RVHe-AHM-gyl. The amino acid sequence and the base sequence of the V region of the H chain contained in this plasmid HEF-RVHe-AHM-gyl are shown in SEQ ID NO: 23.

### 3-2. Construction of the H chain hybrid V region

15                   Two H chain hybrid V regions were constructed. One is a mouse-human hybrid anti-HM1.24 antibody in which the amino acid sequences of FR1 and FR2 are derived from the mouse anti-HM1.24 antibody and those of FR3 and FR4 are from version a of the V region of the H chain of the  
20                   reshaped human anti-HM1.24 antibody, and the other is human-mouse hybrid anti-HM1.24 antibody in which the amino acid sequences of FR1 and FR2 are derived from version a of the V region of the H chain of the reshaped human anti-HM1.24 antibody and those of FR3 and FR4 are  
25                   from the mouse anti-HM1.24 antibody. The amino acid sequences of the CDR regions are all derived from mouse anti-HM1.24 antibody.

                  Two H chain hybrid V regions were constructed by the PCR method. The method is schematically shown in  
30                   Fig. 6 and 7. For the construction of two H chain hybrid V regions, four primers were used. The external primers a (SEQ ID NO: 93) and h (SEQ ID NO: 94) were designed to hybridize with the DNA sequence of the HEF expression vector HEF-VH-gyl. The H chain hybrid construction  
35                   primer HYS (SEQ ID NO: 95) was designed to have the sense

002220 032200 000000 000000

For the construction of the H chain hybrid V region in which the amino acid sequences of FR1 and FR2 are derived from the mouse anti-HM1.24 antibody and those of FR3 and FR4 are from version a of the V region of the H chain of the reshaped human anti-HM1.24 antibody, PCR using the plasmid HEF-1.24H-gyl as a template, the external primer a, and the H chain hybrid primer HYA, and PCR using the plasmid HEF-RVLa-AHM-gyl as a template, the H chain hybrid primer HYS (SEQ ID NO: 95), and the external primer h (SEQ ID NO: 94) were carried out in the first stage of PCR to purify each PCR product. The two PCR products from the first PCR were allowed to assemble by their own complementarity (see International Patent Publication No. WO 92-19759).

For the construction of the H chain hybrid V region in which the amino acid sequences of FR1 and FR2 are derived from version a of the V region of the H chain of the reshaped human anti-HM1.24 antibody and those of FR3 and FR4 are from the mouse anti-HM1.24 antibody, PCR using the plasmid HEF-RVHa-AHM-gyl as a template, the external primer a, and the H chain hybrid primer HYA, and PCR using the plasmid HEF-1.24H-gyl as a template, the H chain hybrid primer HYS, and the external primer h were carried out in the first stage of PCR to purify each PCR product. The two PCR purified products from the first



PCR were allowed to assemble by their own complementarity (see International Patent Publication No. WO 92-19759).

Then, by adding the external primers a and h, a full-length DNA encoding the H chain hybrid V region in which the amino acid sequences of FR1 and FR2 are derived from version a of the V region of the H chain of the reshaped human anti-HM1.24 antibody and those of FR3 and FR4 are from the mouse anti-HM1.24 antibody was amplified in the second PCR stage.

The methods of the first PCR, purification of PCR products, assembling, the second PCR, and cloning into the HEF expression vector HEF-VH-gyl were carried out according to the methods shown in "Example 9. Construction of the V region of the L chain of the reshaped human anti-HM1.24 antibody".

After sequencing the DNA sequence, the plasmid that contains the DNA fragment encoding the correct amino acid sequence of the H chain hybrid V region in which the amino acid sequences of FR1 and FR2 are derived from the mouse anti-HM1.24 antibody and those of FR3 and FR4 are from version a of the V region of the H chain of the reshaped human anti-HM1.24 antibody was termed HEF-MH-RVH-AHM-gyl. The amino acid sequence and the base sequence of the V region of the H chain contained in this plasmid HEF-MH-RVH-AHM-gyl are shown in SEQ ID NO: 97.

Also, the plasmid that contains the DNA fragment encoding the correct amino acid sequence of the H chain hybrid V region in which the amino acid sequences of FR1 and FR2 are derived from version a of the V region of the H chain of the reshaped human anti-HM1.24 antibody and those of FR3 and FR4 are from the mouse anti-HM1.24 antibody was termed HEF-HM-RVH-AHM-gyl. The amino acid sequence and the base sequence of the V region of the H chain contained in this plasmid HEF-HM-RVH-AHM-gyl are shown in SEQ ID NO: 99.

3-3. Construction of versions f to r of the V region of the H chain of the reshaped human anti-HM1.24 antibody

5 Each of versions f, g, h, i, j, k, l, m, n, o, p, q, and r of the V region of the H chain of the reshaped human anti-HM1.24 antibody were constructed as follows.

10 Using as the mutagen primer FS (SEQ ID NO: 102) and FA (SEQ ID NO: 103) designed to mutate threonine at position 75 to serine and valine at position 78 to alanine and as a template DNA the plasmid HEF-RVHe-AHM-gyl by the PCR method, version f was amplified to obtain plasmid HEF-RVHf-AHM-gyl. The amino acid sequence and the base sequence of the V region of the H chain contained in this plasmid HEF-RVHf-AHM-gyl are shown in SEQ ID NO: 25.

15 Using as the mutagen primer GS (SEQ ID NO: 104) and GA (SEQ ID NO: 105) designed to mutate alanine at position 40 to arginine and, as a template DNA, the plasmid HEF-RVHa-AHM-gyl, version g was amplified to obtain plasmid HEF-RVHg-AHM-gyl. The amino acid sequence and the base sequence of the V region of the H chain contained in this plasmid HEF-RVHg-AHM-gyl are shown in SEQ ID NO: 27.

20 Using as the mutagen primer FS (SEQ ID NO: 102) and FA (SEQ ID NO: 103) and, as a template DNA, the plasmid HEF-RVHb-AHM-gyl, version h was amplified to obtain plasmid HEF-RVHh-AHM-gyl. The amino acid sequence and the base sequence of the V region of the H chain contained in this plasmid HEF-RVHh-AHM-gyl are shown in SEQ ID NO: 29.

25 Using as the mutagen primer IS (SEQ ID NO: 106) and IA (SEQ ID NO: 107) designed to mutate arginine at position 83 to alanine and serine at position 84 to

phenylalanine and, as a template DNA, the plasmid  
HEF-RVHh-AHM-gyl, version i was amplified to obtain  
plasmid HEF-RVHi-AHM-gyl. The amino acid sequence and  
the base sequence of the V region of the H chain  
5 contained in this plasmid HEF-RVHi-AHM-gyl are shown in  
SEQ ID NO: 31.

Using as the mutagen primer JS (SEQ ID NO: 108)  
and JA (SEQ ID NO: 109) designed to mutate arginine at  
position 66 to lysine and, as a template DNA, the plasmid  
10 HEF-RVHf-AHM-gyl, version j was amplified to obtain  
plasmid HEF-RVHj-AHM-gyl. The amino acid sequence and  
the base sequence of the V region of the H chain  
contained in this plasmid HEF-RVHj-AHM-gyl are shown in  
SEQ ID NO: 33.

Using as the mutagen primer KS (SEQ ID NO: 110)  
and KA (SEQ ID NO: 111) designed to mutate glutamic acid  
at position 81 to glutamine and, as a template DNA, the  
plasmid HEF-RVHh-AHM-gyl, version k was amplified to  
obtain plasmid HEF-RVHk-AHM-gyl. The amino acid sequence  
20 and the base sequence of the V region of the H chain  
contained in this plasmid HEF-RVHk-AHM-gyl are shown in  
SEQ ID NO: 35.

Using as the mutagen primer LS (SEQ ID NO: 112)  
and LA (SEQ ID NO: 113) designed to mutate glutamic acid  
25 at position 81 to glutamine and serine at position 82B to  
isoleucine and, as a template DNA, the plasmid  
HEF-RVHh-AHM-gyl, version l was amplified to obtain  
plasmid HEF-RVHl-AHM-gyl. The amino acid sequence and  
the base sequence of the V region of the H chain  
30 contained in this plasmid HEF-RVHl-AHM-gyl are shown in  
SEQ ID NO: 37.

Using as the mutagen primer MS (SEQ ID NO: 114)  
and MA (SEQ ID NO: 115) designed to mutate glutamic acid  
at position 81 to glutamine, serine at position 82b to

002222 003333 004444 005555 006666 007777 008888 009999

isoleucine, and threonine at position 87 to serine and, as a template DNA, the plasmid HEF-RVHh-AHM-gyl, version m was amplified to obtain plasmid HEF-RVHm-AHM-gyl. The amino acid sequence and the base sequence of the V region of the H chain contained in this plasmid HEF-RVHm-AHM-gyl are shown in SEQ ID NO: 39.

Using as the mutagen primer NS (SEQ ID NO: 116) and NA (SEQ ID NO: 117) designed to mutate serine at position 82B to isoleucine and, as a template DNA, the plasmid HEF-RVHh-AHM-gyl, version n was amplified to obtain plasmid HEF-RVHn-AHM-gyl. The amino acid sequence and the base sequence of the V region of the H chain contained in this plasmid HEF-RVHn-AHM-gyl are shown in SEQ ID NO: 41.

Using as the mutagen primer OS (SEQ ID NO: 118) and OA (SEQ ID NO: 119) designed to mutate threonine at position 87 to serine and, as a template DNA, the plasmid HEF-RVHh-AHM-gyl, version o was amplified to obtain plasmid HEF-RVHo-AHM-gyl. The amino acid sequence and the base sequence of the V region of the H chain contained in this plasmid HEF-RVHo-AHM-gyl are shown in SEQ ID NO: 43.

Using as the mutagen primer PS (SEQ ID NO: 120) and PA (SEQ ID NO: 121) designed to mutate valine at position 78 to alanine and, as a template DNA, the plasmid HEF-RVHa-AHM-gyl, version p was amplified by the PCR method to obtain plasmid HEF-RVHp-AHM-gyl. The amino acid sequence and the base sequence of the V region of the H chain contained in this plasmid HEF-RVHp-AHM-gyl are shown in SEQ ID NO: 45.

Using as the mutagen primer QS (SEQ ID NO: 122) and QA (SEQ ID NO: 123) designed to mutate threonine at position 75 to serine and, as a template DNA, the plasmid HEF-RVHa-AHM-gyl, version q was amplified by the PCR

5 Using as the mutagen primer CS (SEQ ID NO: 87)  
and CA (SEQ ID NO: 88) and, as a template DNA, the  
plasmid HEF-RVHp-AHM-gyl, version r was amplified by the  
PCR method to obtain plasmid HEF-RVHr-AHM-gyl. The amino  
acid sequence and the base sequence of the V region of  
10 the H chain contained in this plasmid HEF-RVHr-AHM-gyl  
are shown in SEQ ID NO: 49.

The Escherichia coli that contain each of the  
20 plasmids pUC19-RVLa-AHM-gk and pUC19-RVHr-AHM-gyl was  
termed Escherichia coli DH5 $\alpha$  (pUC19-RVLa-AHM-gk) and  
Escherichia coli DH5 $\alpha$  (pUC19-RVHr-AHM-gyl), respectively,  
and have been internationally deposited on August 29,  
1996, with the National Institute of Bioscience and  
25 Human-Technology, Agency of Industrial Science and  
Technology, MITI (Higashi 1-Chome 1-3, Tsukuba city,  
Ibalaki prefecture, Japan) under the accession numbers  
FERM BP-5645 and FERM BP-5643, respectively, under the  
provisions of the Budapest Treaty.

In order to evaluate each chain of the reshaped human anti-HM1.24 antibody, the reshaped human

anti-HM1.24 antibody and the chimera anti-HM1.24 antibody as a positive control antibody were allowed to express. In constructing each of version b and after of the V region of the H chain of the reshaped human anti-HM1.24 antibody, the H chain hybrid antibody was allowed to express in order to investigate which amino acid sequence in the FR should be substituted. Furthermore, it was expressed in combination with the chimera H chain in order to evaluate version a of L chain of the reshaped human anti-HM1.24 antibody.

#### 4-1. Expression of the reshaped human anti-HM1.24 antibody

Ten  $\mu$ g each of the expression vector (HEF-RVHa-AHM-gyl to HEF-RVHr-AHM-gyl) for the H chain of the reshaped human anti-HM1.24 antibody and the expression vector (HEF-RVLb-AHM-gk or HEF-RVLb-AHM-gk) for the L chain of the reshaped human anti-HM1.24 antibody were cotransformed into COS-7 cells by electroporation using the Gene Pulser instrument (manufactured by BioRad). Each DNA (10  $\mu$ g) was added to 0.8 ml aliquots of  $1 \times 10^7$  cells/ml in PBS, and was subjected to pulses at 1500 V and a capacity of 25  $\mu$ F.

After the recovery period of 10 minutes at room temperature, the electroporated cells were added to 30 ml of DHEM culture liquid (manufactured by GIBCO) containing 10%  $\gamma$ -globulin-free bovine fetal serum. After incubation of 72 hours in the CO<sub>2</sub> incubator BNA120D (manufactured by TABAI) under the condition of 37°C and 5% CO<sub>2</sub>, the culture supernatant was collected, the cell debris was removed by centrifugation at 1000 rpm for 5 minutes in a centrifuge 15PR-22 (manufactured by HITACHI) equipped with a centrifuge rotor 03 (manufactured by HITACHI), and a microconcentrator (Centricon 100, manufactured by Amicon) was ultrafiltrated using a centrifuge J2-21 (manufactured by BECKMAN) equipped with a centrifuge

rotor JA-20.1 (manufactured by BECKMAN) at a condition of 2000 rpm, and was used for Cell-ELISA.

4-2. Expression of the chimera anti-HM1.24 antibody

Using ten  $\mu$ g each of the expression vector

5 HEF-1.24H-gyl for the H chain of the chimera human  
anti-HM1.24 antibody and the expression vector  
HEF-1.24L-gk for the L chain of the chimera human  
anti-HM1.24 antibody, the chimera anti-HM1.24 antibody to  
be used for Cell-ELISA was prepared according to the  
10 above-mentioned method for expression of the reshaped  
human anti-HM1.24 antibody.

4-3. Expression of the anti-HM1.24 antibody  
comprising version a of the humanized L chain and  
the chimera H chain

15 Using ten  $\mu$ g each of the expression vector  
HEF-1.24H-gyl for the H chain of the chimera human  
anti-HM1.24 antibody and the expression vector  
HEF-RVLa-AHM-Gk for version a of the L chain of the  
reshaped human anti-HM1.24 antibody, the anti-HM1.24  
20 antibody comprising version a of the humanized L chain  
and the chimera H chain to be used for Cell-ELISA was  
prepared according to the above-mentioned method for  
expression of the reshaped human anti-HM1.24 antibody.

4-4. Expression of the H chain hybrid antibody

25 Using ten  $\mu$ g each of the expression vector  
(HEF-MH-RVH-AHM-gyl or HEF-HM-RVH-AHM-gyl) for the V  
region of the H chain hybrid and the expression vector  
HEF-RVLa-AHM-gk for the L chain of the reshaped human  
anti-HM1.24 antibody, the H chain hybrid antibody to be  
30 used for Cell-ELISA was prepared according to the  
above-mentioned method for expression of the reshaped  
human anti-HM1.24 antibody.

4-5. Measurement of antibody concentration

Concentration of the antibody obtained was

002250 2606060

measured by ELISA. Each well of a 96-well ELISA plate (Maxisorp, manufactured by NUNC) was immobilized by adding 100  $\mu$ l of goat anti-human IgG antibody (manufactured by BIO SOURCE) prepared to a concentration of 1  $\mu$ g/ml with the coating buffer (0.1 M  $\text{NaHCO}_3$ , 0.02%  $\text{NaN}_3$ , pH 9.6) and incubating at room temperature for one hour. After blocking with 100  $\mu$ l of the dilution buffer (50 mM Tris-HCl, 1 mM  $\text{MgCl}_2$ , 0.15 M NaCl, 0.05% Tween 20, 0.02%  $\text{NaN}_3$ , 1% bovine serum albumin (BSA), pH 8.1), 100  $\mu$ l each of serial dilutions of the reshaped human anti-HM1.24 antibody, chimera anti-HM1.24 antibody, and the H chain hybrid antibody that were concentrated by ultrafiltration were added to each well and incubated at room temperature for one hour. Then, after washing, 100  $\mu$ l of alkaline phosphatase-labeled goat anti-human IgG antibody (manufactured by DAKO) was added.

After incubating at room temperature for one hour and washing, 100  $\mu$ l of 1  $\mu$ g/ml substrate solution (Sigma104, p-nitrophenyl phosphate, manufactured by SIGMA) dissolved in the substrate buffer (50 mM  $\text{NaHCO}_3$ , 10 mM  $\text{MgCl}_2$ , pH 9.8) was added, and then the absorbance at 405 nm was measured using the MICROPLATE READER Model 3550 (manufactured by Bio Rad). As the standard for the measurement of concentration, human IgG1k (manufactured by The Binding Site) was used.

5. Establishment of the CHO cell line that stably produces the human anti-HM1.24 antibody

5-1. Construction of the expression vector for the H chain of the reshaped human anti-HM1.24 antibody

By digesting plasmid HEF-RVHr-AHM-gyl with the restriction enzymes PvuI and BamHI, an about 2.8 kbp fragment containing the DNA encoding the EF1 promoter and the V region of the H chain of the reshaped human anti-HM1.24 antibody was purified using 1.5% low melting point agarose gel. Then, the above DNA fragment was



inserted into an about 6 kbp fragment that was prepared by digesting the expression vector used for a human H chain expression vector, DHFR- $\Delta$ E-RVh-PM1f (International Patent Publication No. WO 92-19759), containing the DHFR gene and the gene encoding the constant region of a human H chain with PvuI and BamHI to construct an expression vector, DHFR- $\Delta$ E-HEF-RVHr-AHM-gyl, for the H chain of the reshaped anti-HM1.24 antibody.

#### 5-2. Gene introduction into CHO cells

In order to establish a stable production system of the reshaped anti-HM1.24 antibody, the genes of the above-mentioned expression vectors, DHFR- $\Delta$ E-RVHr-AHM-gyl and HEF-RVLa-AHM-gk, that were linearized by digestion with PvuI were simultaneously introduced into the CHO cell DXB-11 by the electroporation method under the condition similar to the above-mentioned one (transfection into the above-mentioned COS-7 cells).

#### 5-3. Gene amplification by MTX

Among the gene-introduced CHO cells, only those CHO cells in which both of L chain and H chain expression vectors have been introduced can survive in the nucleoside-free  $\alpha$ -MEM culture liquid (manufactured by GIBCO-BRL) to which 500  $\mu$ g/ml G418 (manufactured by GIBCO-BRL) and 10% bovine fetal serum were added, and so they were selected. Subsequently, 10 nM MTX (manufactured by Sigma) was added to the above culture liquid. Among the clones that propagated, those that produce the reshaped anti-HM1.24 antibody in large amounts were selected. As a result, clone #1 that exhibits a production efficiency of about 3  $\mu$ g/ml of the reshaped anti-HM1.24 antibody was obtained and termed the reshaped anti-HM1.24 antibody-producing cell line.

#### 5-4. Construction of the reshaped human anti-HM1.24 antibody

The reshaped anti-HM1.24 antibody was constructed in the following method. The above CHO cells that produce the reshaped anti-HM1.24 antibody were cultured for 10 days using as the medium the nucleoside-free  $\alpha$ -MEM culture liquid (manufactured by GIBCO-BRL) to which 500  $\mu$ g/ml G418 (manufactured by GIBCO-BRL) containing 10%  $\gamma$ -globulin-free bovine fetal serum (manufactured by GIBCO-BRL) were added using the CO<sub>2</sub> incubator BNAS120D (manufactured by TABAI) under the condition of 37°C and 5% CO<sub>2</sub>. On day 8 and 10 after starting the culture the culture liquid was recovered, the cell debris was removed by centrifuging for 10 minutes at 2000 rpm using the centrifuge RL-500SP (manufactured by Tomy Seiko) equipped with the TS-9 rotor, and then filter-sterilized using a bottle top filter (manufactured by FALCON) having a membrane with pores of 0.45  $\mu$ m in diameter.

After an equal amount of PBS(-) was added to the culture liquid of the CHO cells that produce the reshaped human anti-HM1.24 antibody, then the reshaped anti-HM1.24 antibody was affinity-purified using the high-speed antibody purification system ConSep LC100 (manufactured by MILLIPORE) and Hyper D Protein A column (manufactured by Nippon Gaishi) using PBS(-) as the absorption/wash buffer and 0.1 M sodium citrate buffer (pH 3) as the elution buffer according to the attached instructions. The eluted fractions were adjusted to about pH 7.4 by immediately adding 1 M Tris-HCl (pH 8.0) and then using the centrifuging ultrafiltration concentrator Centriprep 10 (manufactured by MILLIPORE), concentration and substitution to PBS(-) was carried out and filter-sterilized using a membrane filter MILLEX-GV (manufactured by MILLIPORE) with a pore size of 0.22  $\mu$ m to obtain the purified reshaped human anti-HM1.24 antibody. Antibody concentration was measured by

absorbance at 280 nm and calculated with 1 µg/ml as 1.35 OD.

Reference Example 11. Determination of activity of the reshaped anti-HM1.24 antibody

5           The reshaped anti-HM1.24 antibody was evaluated for the following antigen binding activity and binding inhibition activity.

1. The method of measurement of antigen binding activity and binding inhibition activity

10          1-1. Measurement of antigen binding activity

Antigen binding activity was measured by the Cell-ELISA using WICH cells. Cell-ELISA plates were prepared as described in the above Example 7.1-2.

15           After blocking, 100 µl of serial dilutions of the reshaped human anti-HM1.24 antibody that was obtained from the concentrate of the culture supernatant of COS-7 cells or purified from the culture supernatant of CHO cells was added to each well. After it was incubated for 2 hours at room temperature and washed,  
20          peroxidase-labeled rabbit anti-human IgG antibody (manufactured by DAKO) was added. After incubating for 2 hours at room temperature and washing, the substrate solution was added and incubated. Then the reaction was stopped by adding 50 µl of 6N sulfuric acid, and  
25          absorbance at 490 nm was measured using the MICROPLATE READER Model 3550 (manufactured by Bio-Rad).

1-2. Measurement of binding inhibition activity

30           The binding inhibition activity by the biotin-labeled mouse anti-HM1.24 antibody was measured by the Cell-ELISA using WISH cells. Cell-ELISA plates were prepared as described above. After blocking, 50 µl of serial dilutions of the reshaped human anti-HM1.24 antibody that was obtained from the concentrate of the culture supernatant of COS-7 cells or purified from the culture supernatant of CHO cells was added to each well,  
35          and 50 µl of 2 µg/ml biotin-labeled mouse anti-HM1.24

antibody was added simultaneously. After incubating at room temperature for two hours and washing, peroxidase-labeled streptavidin (manufactured by DAKO) was added. After incubating at room temperature for one hour and then washing, the substrate solution was added and incubated. Then the reaction was stopped by adding 50 µl of 6N sulfuric acid, and absorbance at 490 nm was measured using the MICROPLATE READER Model 3550 (manufactured by Bio-Rad).

2. Evaluation of the reshaped human anti-HM1.24 antibody

2-1. L chain

Version a of the L chain of the reshaped human anti-HM1.24 antibody was evaluated as mentioned above for measurement of antigen binding activity. As shown in Fig. 8, when version a of the L chain is expressed in combination with the chimera H chain it has shown a similar level of antigen binding activity. However, in consideration of further increase in activity and of compatibility with the H chain, version b of the L chain was constructed. Versions a and b of the L chain were evaluated together for antigen binding activity and of binding inhibition activity when combined with versions a, b, f, or h of the H chain. As shown in Fig. 9, 10, 11, and 12, version a of the L chain had a higher activity than version b in both activities in all versions a, b, f, and h of the H chain. Therefore, version a of the L chain of the reshaped human anti-HM1.24 antibody was used for the following experiment.

2-2. H chain versions a to e

Versions a to e of the H chain of the reshaped human anti-HM1.24 antibody were evaluated in combination with the version a of the L chain as mentioned above for measurement of antigen binding activity and for binding inhibition activity. The result, as shown in Fig. 11, 13, 14, and 15, indicated that all versions were weaker

in both activities as compared to the chimera anti-HM1.24 antibody, suggesting that further amino acid substitution is required.

### 2-3. The H chain hybrid antibody

5           The H chain hybrid antibody was evaluated as mentioned above for measurement of antigen binding activity. The result, as shown in Fig. 16, indicated that the human-mouse hybrid anti-HM1.24 antibody has shown a similar activity to that of the chimera  
10 anti-HM1.24 antibody for antigen binding activity, whereas the mouse-human hybrid anti-HM1.24 antibody had a weaker activity than the chimera anti-HM1.24 antibody. This indicated that, in order to construct the reshaped human anti-HM1.24 antibody having the antigen binding  
15 activity similar to that of the chimera anti-HM1.24 antibody, it is necessary to convert amino acids included in FR3 or FR4 among those contained the V region of the H chain.

### 2-4. Versions f to r of the H chain

20           Version f of the H chain of the reshaped human anti-HM1.24 antibody was evaluated as mentioned above for measurement of antigen binding activity. The result, as shown in Fig. 17, indicated that its antigen binding activity is decreased as compared to the chimera  
25 anti-HM1.24 antibody, but is increased as compared to the above versions a to c, suggesting that any of the four amino acids at positions 67, 69, 75, and 78 that were newly converted in this version is responsible for the activity of the reshaped human antibody.

30           Version g of the H chain of the reshaped human anti-HM1.24 antibody was evaluated as mentioned above for measurement of antigen binding activity. The result, as shown in Fig. 18 and 19, indicated that this version has exhibited a similar level of activity to that of the  
35 above version a at most, revealing that, as shown for the above H chain human-mouse hybrid antibody, the amino acid at position 40 that was converted in this version is not

responsible for the increase in the activity of the reshaped human antibody.

5 Versions h to j of the H chain of the reshaped human anti-HM1.24 antibody were evaluated as mentioned above for measurement of antigen binding activity and of binding inhibition activity. The result, as shown in Fig. 20, 21, 22, and 23, indicated that all versions were weaker for both activities as compared to the chimera anti-HM1.24 antibody and were similar to the  
10 above-mentioned f, suggesting that the amino acids at positions 67 and 69 among the four amino acids that were newly converted in version f are not responsible for the increase in the activity of the reshaped human antibody.

15 Versions k to p of the H chain of the reshaped human anti-HM1.24 antibody were evaluated as mentioned above for measurement of antigen binding activity and of binding inhibition activity. The result, as shown in Fig. 24, 25, 26, and 27, indicated that all versions were weaker for both activities as compared to the chimera anti-HM1.24 antibody and were similar to the  
20 above-mentioned h, suggesting that the amino acids at position 80 and after that were newly converted in these six versions are not responsible for the increase in the activity of the reshaped human antibody.

25 Version q of the H chain of the reshaped human anti-HM1.24 antibody was evaluated as mentioned above for measurement of antigen binding activity and of binding inhibition activity. The result, as shown in Fig. 25 and 27, indicated that this version was weaker for both  
30 activities as compared to the above version h or version p and was similar to that of the above-mentioned a at most, suggesting that substitution of the amino acid at position 78 is essential for the increase in the activity of the reshaped human antibody.

35 Version r of the H chain of the reshaped human anti-HM1.24 antibody were evaluated by the method mentioned above. The result, as shown in Fig. 15 and 28,

The above results indicated that the minimum conversion required for the reshaped human anti-HM1.24 antibody to have a similar level of antigen binding activity to that of the mouse anti-HM1.24 antibody or the chimera anti-HM1.24 antibody is the amino acids at positions 30, 71, and 78 and, furthermore, 73.

10                   The antigen binding activity and the binding inhibition activity for H chain versions a to r of the reshaped human anti-HM1.24 antibody are summarized in Table 2.

Table 2

H chain version	Antigen binding activity	Binding inhibition activity
a	+	+
b	+	+
c	+	+
d	+	not measured
e	+	not measured
f	++	++
g	+	+
h	++	++
i	++	++
j	++	++
k	++	++
l	++	++
m	++	++
n	++	++
o	++	++
p	++	++
q	+	+
r	+++	+++

5 Furthermore, the amino acid sequences of the reshaped human anti-HM1.24 antibody and versions a and b of the L chain are shown in Table 3, and those of versions a to r of the H chain of the reshaped human anti-HM1.24 antibody are shown in Tables 4 to 6.



The amino acid sequence of the L chain V region

[illegible]

Table 4

The amino acid sequence of the H chain V region (1)

	FR1																		CDR1					FR2					
	1									2									3					4					
	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27		
AHM	Q	V	Q	L	Q	Q	S	G	A	E	L	A	R	P	G	A	S	V	K	L	S	C	K	A	S	G	Y	T	F
HuSGI	E	V	Q	L	V	Q	S	G	A	D	V	K	K	P	G	X	S	V	X	V	S	C	K	A	S	G	Y	T	F
HG3	Q	V	Q	L	V	Q	S	G	A	E	V	K	K	P	G	A	S	V	K	V	S	C	K	A	S	G	Y	T	F
RVHa	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
RVHb	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
RVHc	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
RVHd	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
RVHe	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
RVHf	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
RVHg	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
RVHh	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
RVHi	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
RVHj	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
RVHk	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
RVHl	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
RVHm	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
RVHn	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
RVHo	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
RVHp	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
RVHq	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	
RVHr	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	-	

002250" 85050550

The amino acid sequence of the H chain V region (2)

[illegible]

Table 6

The amino acid sequence of the H chain V region

	CDR3	FR4
	10	11
	57890ABJK12	34567890123
AHM	GLRRGGYYFDY	WGQGTTLTVSS
HuSGI		WGQGTTLTVSS
JH6		WGQGTTLTVSS
RVHa	-----	-----
RVHb	-----	-----
RVHc	-----	-----
RVHd	-----	-----
RVHe	-----	-----
RVHf	-----	-----
RVHg	-----	-----
RVHh	-----	-----
RVHi	-----	-----
RVHj	-----	-----
RVHk	-----	-----
RVHl	-----	-----
RVHm	-----	-----
RVHn	-----	-----
RVHo	-----	-----
RVHp	-----	-----
RVHq	-----	-----
RVHr	-----	-----

25            3. Evaluation of the purified reshaped human  
anti-HM1.24 antibody

             The purified reshaped human anti-HM1.24  
antibody was evaluated for the above-mentioned antigen  
binding activity and binding inhibition activity. The  
30        result, as shown in Fig. 31 and 32, indicated that the  
reshaped human anti-HM1.24 antibody has a similar level  
of antigen binding activity and binding inhibition  
activity to that of the chimera anti-HM1.24 antibody.  
This fact indicated that the reshaped human anti-HM1.24  
35        antibody has the same antigen binding activity as the  
mouse anti-HM1.24 antibody.

Reference example 12. Construction of the hybridoma  
that produces the mouse anti-HM1.24 monoclonal  
antibody

5 The hybridoma that produces the mouse anti-HM1.24  
monoclonal antibody was prepared according to the method  
described in Goto, T. et al., Blood (1994) 84, 1992-1930.

The Epstein-Barr virus nuclear antigen  
(EBNA)-negative plasma cell line KPC-32 ( $1 \times 10^7$  cells)  
derived from the bone marrow of human patients with  
10 multiple myeloma (Goto, T. et al., Jpn. J. Clin. Hematol.  
(1991) 32, 1400) was intraperitoneally given twice to  
BALB/c mice (manufactured by Charles River) every six  
weeks.

In order to further elevate the titer of antibody  
15 production,  $1.5 \times 10^6$  KPC-32 cells were injected into the  
spleen of the mice three days before sacrificing the  
animals (Goto, T. et al., Tokushima J. Exp. Med. (1990)  
37, 89). After sacrificing the mice, the spleen was  
removed, and the spleen cells removed according to the  
20 method of Groth, de St. & Schreidegger (Cancer Research  
(1981) 41, 3465) were subjected to cell fusion with the  
myeloma cells SP2/0.

Antibody in the supernatant of the hybridoma culture  
was screened by the ELISA (Posner, M.R. et al., J.  
25 Immunol. Methods (1982) 48, 23) using the KPC-32  
cell-coated plates.  $5 \times 10^4$  KPC-32 cells were suspended  
in 50 ml of PBS and dispensed into 96-well plates  
(U-bottomed, Corning, manufactured by Iwaki). After  
blocking with PBS containing 1% bovine serum albumin  
30 (BSA), the supernatant of the hybridoma was added and  
incubated at  $4^\circ\text{C}$  for 2 hours. Subsequently,  
peroxidase-labeled anti-mouse IgG goat antibody  
(manufactured by Zymed) was reacted at  $4^\circ\text{C}$  for 1 hour,  
washed once, and was reacted with the o-phenylenediamine  
35 substrate solution (manufactured by Sumitomo Bakelite) at  
room temperature for 30 minutes.

After stopping the reaction with 2N sulfuric acid,

absorbance at 492 nm was measured using the ELISA reader (manufactured by Bio-Rad). In order to remove the hybridoma that produces antibody against human immunoglobulin, the positive hybridoma culture supernatant had previously been adsorbed to human serum, and the reactivity to other sub-cellular components was screened. Positive hybridomas were selected and their reactivity to various cell lines and human samples was investigated using flow cytometry. The finally selected hybridoma clones were cloned twice, were injected into the abdominal cavity of the pristane-treated BALB/c mice and then the ascitic fluid was obtained therefrom.

Monoclonal antibody was purified from the mouse ascites by ammonium sulfate precipitation and Protein A affinity chromatography kit (Ampure PA, manufactured by Amersham). The purified antibody was conjugated to fluorescein isocyanate (FITC) using the Quick Tag FITC conjugation kit (manufactured by Boehringer Mannheim).

As a result, the monoclonal antibody produced by 30 hybridoma clones reacted with KPC-32 and RPMI 8226 cells. After cloning, the reactivity of the supernatant of these hybridomas with other cell lines and peripheral blood-derived monocytes was investigated.

Of them, three clones were monoclonal antibodies that specifically react with plasma cells. Out of these three clones, the hybridoma clone having the clone that is most useful for flow cytometry analysis and that has complement-dependent cytotoxicity was selected and termed HM1.24. The subclass of monoclonal antibody produced by this hybridoma was determined by ELISA using subclass-specific anti-mouse rabbit antibody (manufactured by Zymed). Anti-HM1.24 antibody had a subclass of IgG2a  $\kappa$ . The hybridoma that produces the anti-HM1.24 antibody was internationally deposited on September 14, 1995, with the National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology, MITI (Higashi 1-Chome 1-3,

002250 00000000

Tsukuba city, Ibaraki prefecture, Japan) under the accession number FERM BP-5233 under the provisions of the Budapest Treaty.

Reference example 13. Cloning of cDNA encoding the HM1.24 antigen polypeptide

1. Construction of cDNA library

1) Preparation of total RNA

The cDNA that encodes the HM1.24 antigen which is an antigen polypeptide specifically recognized by mouse monoclonal antibody HM1.24 was isolated as follows.

From the human multiple myeloma cell line KPMM2, total RNA was prepared according to the method of Chirgwin et al. (Biochemistry, 18, 5294 (1979)). Thus,  $2.2 \times 10^8$  KPMM2 cells were completely homogenized in 20 ml of 4 M guanidine isocyanate (manufactured by Nacalai Tesque Inc.).

The homogenate was layered on the 5.3 M cesium chloride layer in the centrifuge tube, which was then centrifuged using Beckman SW40 rotor at 31,000 rpm at 20 °C for 24 hours to precipitate RNA. The RNA precipitate was washed with 70% ethanol, and dissolved in 300 µl of 10 mM Tris-HCl (pH 7.4) containing 1 mM EDTA and 0.5% SDS. After adding Pronase (manufactured by Boehringer) thereto to a concentration of 0.5 mg/ml, it was incubated at 37 °C for 30 minutes. The mixture was extracted with phenol and chloroform to precipitate RNA. Then, the RNA precipitate was dissolved in 200 µl of 10 mM Tris-HCl (pH 7.4) containing 1 mM EDTA.

2) Preparation of poly(A)+RNA

Using about 500 µg of the total RNA prepared as above as a raw material, poly(A)+RNA was purified using the Fast Track 2.0m RNA Isolation Kit (manufactured by Invitrogen) according to the instructions attached to the kit.

3) Construction of cDNA library

Using 10 µg of the above poly(A)+RNA as a raw

material, double strand cDNA was synthesized using the  
cDNA synthesizing kit TimeSaver cDNA Synthesis Kit  
(manufactured by Pharmacia) according to the instructions  
attached to the kit and, using the Directional Cloning  
5 Toolbox (manufactured by Pharmacia), EcoRI adapter was  
linked thereto according to the instructions attached to  
the kit. Kination and restriction enzyme NotI treatment  
of the EcoRI adapter were carried out according to the  
instructions attached to the kit. Furthermore, the  
10 adapter-attached double strand cDNA having a size of  
about 500 bp or higher was isolated and purified using  
1.5% agarose gel (manufactured by SIGMA) to obtain about  
40 µl of adapter-attached double strand cDNA.

The adapter-attached double strand cDNA thus  
15 prepared was linked using pCOS1 vector (Japanese  
Unexamined Patent Publication (Kokai) No. 8(1996)-255196)  
and T4 DNA ligase (manufactured by GIBCO BRL) that had  
previously been treated with restriction enzymes EcoRI  
and NotI and alkaline phosphatase (manufactured by Takara  
20 Shuzo) to construct a cDNA library. The constructed cDNA  
library was transduced into Escherichia coli strain DH5  
(manufactured by GIBCO BRL) and the total size was  
estimated to be about  $2.5 \times 10^6$  independent cells.

## 2. Cloning by direct expression

### 25 1) Transfection into COS-7 cells

cDNA was amplified by culturing about  $5 \times 10^5$  clones  
of the above transduced Escherichia coli in the 2-YT  
medium (Molecular Cloning: A Laboratory Manual, Sambrook  
et al., Cold Spring Harbor Laboratory Press, (1989))  
30 containing 50 µg/ml of ampicillin, and plasmid DNA was  
recovered from the Escherichia coli by the alkali method  
(Molecular Cloning: A Laboratory Manual, Sambrook et al.,  
Cold Spring Harbor Laboratory Press, (1989)). The  
plasmid DNA obtained was transfected into COS-7 cells by  
35 electroporation using the Gene Pulser instrument  
(manufactured by BioRad).



Thus, 10 µg of the purified plasmid DNA was added to 0.8 ml of COS-7 cells that were suspended into PBS at a concentration of  $1 \times 10^7$  cells/ml, and was subjected to pulses at 1500 V and a capacity of 25 µF. After 10 minutes of recovery period at room temperature, the electroporated cells were cultured in the DMEM medium (manufactured by GIBCO BRL) supplemented with 10% bovine fetal serum under the condition of 37 °C and 5% CO<sub>2</sub> for three days.

2) Preparation of the panning dish

A panning dish coated with the mouse anti-HM1.24 antibody was prepared by the method of B. Seed et al. (Proc. Natl. Acad. Sci. USA, 84, 3365-3369 (1987)).

Thus, the mouse anti-HM1.24 antibody was added to 50 mM Tris-HCl, pH 9.5, to a concentration of 10 µg/ml. Three ml of the antibody solution thus prepared was added to a tissue culture plate with a diameter of 60 mm and incubated at room temperature for 2 hours. After washing three times with PBS containing 0.15 M NaCl, 5% bovine fetal serum, 1 mM EDTA, and 0.02% NaN<sub>3</sub> was added, and after blocking, it was used for the following cloning.

3) Cloning of cDNA library

The COS-7 cells transfected as described above were detached by PBS containing 5 mM EDTA, and then washed once with PBS containing 5% bovine fetal serum. It was then suspended in PBS containing 5% bovine fetal serum and 0.02% NaN<sub>3</sub> to a concentration of about  $1 \times 10^6$  cells/ml, which was added to the panning dish prepared as above and incubated at room temperature for 2 hours. After washing three times with PBS containing 5% bovine fetal serum and 0.02% NaN<sub>3</sub>, plasmid DNA was recovered from the cells bound to the panning dish using a solution containing 0.6% SDS and 10 mM EDTA.

The recovered plasmid DNA was transduced again to Escherichia coli DH5α. After amplifying the plasmid DNA as above, it was recovered by the alkali method. The

recovered plasmid DNA was transfected into COS-7 cells by the electroporation method to recover plasmid DNA from the bound cells as described above. The same procedure was repeated one more time, and the recovered plasmid DNA was digested with restriction enzymes EcoRI and NotI. As a result, concentration of the insert with a size of about 0.9 kbp was confirmed. Fifty µg of Escherichia coli transduced with part of the recovered plasmid DNA was inoculated to the 2-YT agar plate containing 50 µg/ml of ampicillin. After culturing overnight, plasmid DNA containing a single colony was recovered. It was digested with restriction enzymes EcoRI and NotI and clone p3.19 having an insert of 0.9 kbp was obtained.

The base sequence of this clone was determined by reacting using PRISM, Terminator Cycle Sequencing kit (manufactured by Perkin Elmer) according to the instructions attached to the kit. The amino acid sequence and the base sequence thereof are shown in SEQ ID NO: 128.

The cDNA encoding the polypeptide having the amino acid sequence as set forth in SEQ ID NO: 128 was inserted into the XbaI cleavage site of pUC19 vector, and has been prepared as plasmid pRS38-pUC19. The Escherichia coli that contains this plasmid pRS38-pUC19 has been internationally deposited on October 5, 1993, as Escherichia coli DH5α (pRS38-pUC19), with the National Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology, MITI (Higashi 1-Chome 1-3, Tsukuba city, Ibaraki prefecture, Japan) under the accession number FERM BP-4434 under the provisions of the Budapest Treaty (see Japanese Unexamined Patent Publication (Kokai) No. 7(1995)-196694).

#### EXAMPLES

As an example of natural humanized antibodies composed of the natural FR sequences of the present

Example 1.

5        Mouse monoclonal anti-HM1.24 antibody was humanized  
as the reshaped human anti-HM1.24 antibody by CDR-  
grafting as described in Reference Examples. Each FR of  
human antibody HG3 for FR1 to FR3 and the FR4 of human  
antibody JH6 for FR4 were selected for the construction  
10 of the humanized H chain. The result on the study of the  
FR amino acid residues indicated that amino acid  
substitution was required at four sites (FR1/30, FR3/71,  
73, 78) (Tables 7 and 8). This humanized antibody had an  
antigen binding activity similar to that of the original  
15 antibody. This humanized antibody (humanized antibody  
comprising RVL<sub>a</sub>/RVH<sub>r</sub>) was used as the primary design  
antibody.

A) L chain

FR1	1	2	CDR1	3	FR2	4	CDR2	5
12345678901234567890123			45678901234		567890123456789		0123456	
DIVMTQSHKFMSTSVGDRVITC			KASQDVNTAVA		WYQQKPGQSPKLLIY		SASNRYT	
DIQMTQSPSSLSASVGDRVITC					WYQQKPGKAPKLLIY			
DIQMTQSPSSLSASVGDRVITC					WYQQKPGKAPKLLIY			

## Primary design (RVLa

## Secondary design

FR3	6	7	8	9	CDR3	10	FR4
78901234567890123456789012345678					901234567		8901234567
GVPDRITGSGSGTDFFTISSVQAEDALAYC					QQHYSTPFT		FGSGTKLEIK
GVPSPRFSGSGSGTDFTLTISLQPEDFATYYC							FGQGTVKEIK
GVPSPRFSGSGSGTDFFTISLQPEDIATYYC							FGQGTVKEIK

Primary design (RVLa)

## Secondary design

Design of V region of Natural Humanized Antibody

B) H chain

	FR1	1	2	3	CDR1	FR2	4	5	CDR2	6
HM1.24	123456789012345678901234567890				12345	67890123456789			012A3456789012345	
HuSGI	QVQLQQSCAELARPGASVKLSCKASGYTFT				PYWMQ	WVKRPGQGLEWIG			SIFPGDTRYSQKFKG	
HG3	EVQLVQSCADVKKPGXSVXVSKASGYTFS					WVRQAPGXGLDWVG				
	QVQLVQSGAEVKKPGASVKVSKASGYTFN					WVRQAPGQGLEWMG				

Primary design (RVHr)

Secondary design (2ndRVH)

HM1.24  
HuSGI  
HG3/JH6

	FR3	7	8	9	CDR3	10	11	FR4
67890123456789012ABC345678901234					57890ABJK12			34567890123
KATLTADKSSSTAYMQLSILAFEDSAVYYCAR					GLRRGGYFPDY			WGQGTTLTVSS
RVXTXDXSXNTAYMELSSLRSEDVAVYYCAR								WGQGTTLTVSS
RVWTMRDTSTSTVYMESSLRSEDVAVYYCAR								WGQGTTLTVSS

Primary design (RVHr)

Secondary design (2ndRVH)

-----A-K-----A-----

-----I-A-K-----A-----

(1) The construction of H chain

For the FR of the primary design antibody, homology search on human FRs found in nature was carried out using such databases as SeissPlot, GenBank, PRF, PIR, and GenPept. First, 50 human FRs were found that have completely matching amino acid sequences for FR1. Thus, the FR1 of the primary design antibody already had a natural sequence. Since no amino acid substitution has been made for FR2 and FR4, 50 and 100 natural FRs including HG3 and JH6 respectively of natural human body were found.

On the other hand, no complete matches were found for FR3. As the FR3 that had the highest homology, S46463 having a homology of 96.875%, 1921296C, HUMIGHRF 1, U00583 1 and the like were found (symbols are all accession numbers for the database).

Thus, in the primary design antibody, FR3 was the FR containing artificial amino acid residues that are not found in nature. The amino acid sequence is compared with that of the human antibody S46463 that had the highest homology in Table 9.

FR3 of primary design antibody

```

      10      20      30
      RYTMADKSTSTAYMELSSLRSEDVAVYYCAR
      .....
      .....
      VRQAPGQGLEWMGRIIPILGIANYAQKFQGRVTITADKSTSTAYMELSSLRSEDVAVYYCAR
      40      50      60      70      80      90
```

FR3 of S46463 antibody

30

The amino acid residue at position 70 was methionine in the FR3 of the primary design antibody and was isoleucine in the FR3 of the human antibody S46463. The other amino acid sequences have shown complete matches. Thus, the amino acid residue at position 70 in the primary design antibody was replaced with isoleucine to convert it to a naturally occurring FR3. Accordingly,

35

the secondary design antibody obtained is a CDR-grafting antibody comprising the natural human FR of the human antibody S46463. The secondary design antibody thus constructed comprises FRs that are all found in nature.

5           (2) Construction of the H chain V region of natural humanized anti-HM1.24 antibody

          The H chain V region of the natural humanized anti-HM 1.24 antibody was constructed by mutagenesis using PCR. The mutagen primers SS (SEQ ID NO: 124) and  
10       SA (SEQ ID NO: 125) were designed to mutate methionine at position 69 to isoleucine.

          After the above primer was amplified using plasmid HEF-RVHr-AHM-gyl as a template, the final product was purified, digested with BamHI and HindIII, and the DNA  
15       fragment obtained was cloned into an expression vector HEF-VH-gyl to obtain a plasmid HEF-RVHs-AHM-gyl. The amino acid sequence and the nucleotide sequence of the V region of the H chain contained in this plasmid HEF-RVHs-AHM-gyl are shown in SEQ ID NO: 126.

20           The region encoding the variable region of the above-mentioned plasmid HEF-RVHs-AHM-gyl was digested with restriction enzymes HindIII and BamHI to make a restriction fragment. This was inserted into the BamHI and HindIII sites of plasmid vector pUC19. The plasmid  
25       obtained was termed pUC19-RVHs-AHM-gyl.

Escherichia coli that contains pUC19-RVHs-AHM-gyl was designated as Escherichia coli DH5α (pUC19-RVHs-AHM-gyl) and has been internationally deposited on September 29, 1997, with the National  
30       Institute of Bioscience and Human-Technology, Agency of Industrial Science and Technology, MITI (Higashi 1-Chome 1-3, Tsukuba city, Ibaraki prefecture, Japan) under the accession number FERM BP-6127 under the provisions of the Budapest Treaty.

35           2) Analysis of L chain

Although amino acids of the FRs were not substituted in the construction of the L chain of the primary design antibody, homology search was conducted also for these FRs, since the human antibody REI used was a Reshaped FR (Riechmann, L. et al., Nature (1988) 332, 323-327) that had already been subjected to amino acid substitution. The result confirmed the presence of natural sequences corresponding to the reshaped FRs. Thus, it was demonstrated that no amino acid substitution is required for FRs of L chain.

Example 2. Production of natural humanized anti-HM1.24 antibody

(1) Expression of natural humanized anti-HM1.24 antibody

Ten µg each of the expression vector (HEF-RVHs-AHM-gyl) for H chain of natural humanized anti-HM1.24 antibody and the expression vector (HEF-RVLa-AHM-gk) for L chain of reshaped human anti-HM1.24 antibody was cotransformed into COS cells by electroporation using the Gene Pulser instrument (manufactured by BioRad). Each DNA (10 µg) was added to 0.8 ml aliquots of  $1 \times 10^7$  cells/ml in PBS, and was subjected to pulses at 1500 V and a capacity of 25 µF.

After a recovery period of 10 minutes at room temperature, the electroporated cells were added to 30 ml of DHEM culture liquid (manufactured by GIBCO) containing 10% γ-globulin-free bovine fetal serum. After incubation of 72 hours in a CO<sub>2</sub>-incubator BNA120D (manufactured by TABAI) under the condition of 37°C and 5% CO<sub>2</sub>, the culture supernatant was collected, and the cell debris was removed by centrifugation at 1000 rpm for 5 minutes in a centrifuge 505PR-22 (manufactured by HITACHI) equipped with a centrifuge rotor 03 (manufactured by HITACHI). Then ultrafiltration was carried out with a microconcentrator (Centricon 100, manufactured by Amicon)



using a centrifuge J2-21 (manufactured by BECKMAN)  
equipped with a centrifuge rotor JA-20.1 (manufactured by  
BECKMAN), at a condition of 2000 rpm, and filter-  
sterilization was carried out using a filter Milex GV13mm  
5 (manufactured by Millipore) to obtain a product which was  
used for Cell-ELISA.

(2) Measurement of antibody concentration

Concentration of the antibody obtained was  
measured by ELISA. To each well of a 96-well ELISA plate  
10 (Maxisorp, manufactured by NUNC) was added 100 µl of goat  
anti-human IgG antibody (manufactured by BIO SOURCE)  
prepared to a concentration of 1 µg/ml with the coating  
buffer (0.1 M NaHCO<sub>3</sub>, 0.02% NaN<sub>3</sub>, pH 9.6) and the plate  
was incubated at room temperature for one hour. After  
15 blocking with 100 µl of the dilution buffer (50 mM  
Tris-HCl, 1 mM MgCl<sub>2</sub>, 0.15 M NaCl, 0.05% Tween 20, 0.02%  
NaN<sub>3</sub>, 1% bovine serum albumin (BSA), pH 8.1), 100 µl each  
of serial dilutions of the natural humanized anti-HM1.24  
antibody was added to each well and the plate was  
20 incubated at room temperature for one hour. Then after  
washing, 100 µl of alkaline phosphatase-labeled goat  
anti-human IgG antibody (manufactured by DAKO) was added.

After incubating at room temperature for one  
hour and washing, 100 µl of 1 mg/ml substrate solution  
25 (Sigma 104, p-nitrophenyl phosphate, manufactured by  
SIGMA) dissolved in substrate buffer (50 mM NaHCO<sub>3</sub>, 10 mM  
MgCl<sub>2</sub>, pH 9.8) was added, and then the absorbance at 405  
nm was measured using the MICROPLATE READER Model 3550  
(manufactured by Bio Rad). As a standard for measurement  
30 of concentration, human IgG1k (manufactured by The  
Binding Site) was used.

(3) Establishment of the CHO cell line that stably  
produces the natural humanized anti-HM1.24 antibody

The CHO cell line that stably produces the  
35 natural humanized anti-HM1.24 antibody can be established

according to the following method.

(3)-1. Construction of an expression vector for an H chain of a natural humanized anti-HM1.24 antibody

By digesting plasmid HEF-RVHS-AHM-gyl with  
5 restriction enzymes PvuI and BamHI, an about 2.8 kbp  
fragment containing DNA encoding an EF1 promoter and a V  
region of the H chain of natural humanized anti-HM1.24  
antibody was purified using 1.5% low melting point  
agarose gel. Then, the above DNA fragment is inserted  
10 into an about 6 kbp fragment that was prepared by  
digesting with PvuI and BamHI the expression vector used  
for a human H chain expression vector, DHFR-ΔE-RVh-PM1f  
(International Patent Publication No. WO 92-19759),  
containing a DHFR gene and a gene encoding a constant  
15 region of a human H chain, so as to construct an  
expression vector, DHFR-ΔE-HEF-RVHS-AHM-gyl, for the H  
chain of the natural humanized anti-HM1.24 antibody.

(3)-2. Gene introduction into CHO cells

In order to establish a stable production  
20 system of the natural humanized anti-HM1.24 antibody, the  
genes of the above-mentioned expression vectors,  
DHFR-ΔE-RVHS-AHM-gyl and HEF-RVLa-AHM-gk, that were  
linearized by digestion with PvuI, were simultaneously  
introduced into the CHO cell DXB-11 by the  
25 electroporation method under the condition similar to the  
above-mentioned one (transfection into the  
above-mentioned COS-7 cells).

(3)-3. Gene amplification by MTX

Of the gene-introduced CHO cells, only those  
30 CHO cells in which both of L chain and H chain expression  
vectors have been introduced can survive in the  
nucleoside-free α-MEM culture liquid (manufactured by  
GIBCO-BRL) to which 500 μg/ml G418 (manufactured by  
GIBCO-BRL) and 10% bovine fetal serum were added, and so  
35 they were selected. Subsequently, 10 nM MTX

(manufactured by Sigma) is added to the above culture. Of the clones that propagated, those that produce a natural humanized anti-HM1.24 antibody in large amount were selected.

5           (3)-4. Construction of the natural humanized  
            anti-HM1.24 antibody

            The natural humanized anti-HM1.24 antibody was produced in the following method. The above CHO cells that produce the natural humanized anti-HM1.24 antibody  
10           were cultured for 10 days using a nucleoside-free  $\alpha$ -MEM culture medium (manufactured by GIBCO-BRL) to which 500  $\mu$ g/ml G418 (manufactured by GIBCO-BRL) containing 10%  $\gamma$ -globulin-free bovine fetal serum (manufactured by GIBCO-BRL) had been added, using a CO<sub>2</sub> incubator BNAS120D  
15           (manufactured by TABAI) under the condition of 37°C and 5% CO<sub>2</sub>. On day 8 and 10 after starting the culture the culture medium was recovered, the cell debris was removed by centrifuging for 10 minutes at 2000 rpm using the centrifuge RL-500SP (manufactured by Tomy Seiko) equipped  
20           with the TS-9 rotor, and then filter-sterilized using a bottle top filter (manufactured by FALCON) having a membrane with pores of 0.45  $\mu$ m in diameter.

            After an equal amount of PBS(-) was added to the culture liquid of the CHO cells that produce the  
25           natural humanized anti-HM1.24 antibody, then the natural humanized anti-HM1.24 antibody was affinity-purified using the high-speed antibody purification system ConSep LC100 (manufactured by MILLIPORE) and Hyper D Protein A column (manufactured by Nippon Gaishi) using PBS(-) as an  
30           absorption buffer and 0.1 M sodium citrate buffer (pH 3) as an elution buffer, according to the attached instructions. The eluted fractions were adjusted to about pH 7.4 by immediately adding 1 M Tris-HCl (pH 8.0) and then using the centrifuging ultrafiltration  
35           concentrator Centriprep 10 (manufactured by MILLIPORE), concentration and substitution to PBS(-) were carried out

and the product was filter-sterilized using a membrane filter MILLEX-GV (manufactured by MILLIPORE) with a pore size of 0.22  $\mu$ m to obtain the purified natural humanized anti-HM1.24 antibody. Concentration of purified antibody was measured by absorbance at 280 nm and calculated as 1  $\mu$ g/ml per 1.35 OD.

Example 3. Determination of activity of the natural humanized anti-HM1.24 antibody

The natural humanized anti-HM1.24 antibody was evaluated for the following antigen binding activity, binding inhibition activity, and ADCC activity.

(1) The method of measurement of antigen binding activity and binding inhibition activity

(1)-1. Measurement of antigen binding activity

Antigen binding activity was measured by Cell-ELISA using WICH cells. Cell-ELISA plates were prepared as described in the above Reference Example 7.1-2.

After blocking, 100  $\mu$ l of serial dilutions of the natural humanized anti-HM1.24 antibody that was obtained from a concentrate of a culture supernatant of COS-7 cells was added to each well. After it was incubated for 2 hours at room temperature and washed, peroxidase-labeled rabbit anti-human IgG antibody (manufactured by DAKO) was added. After incubating for 2 hours at room temperature and washing, a substrate solution was added and incubated. Then the reaction was stopped by adding 50  $\mu$ l of 6N sulfuric acid, and absorbance at 490 nm was measured using the MICROPLATE READER Model 3550 (manufactured by Bio-Rad).

(1)-2. Measurement of binding inhibition activity

The binding inhibition activity by the biotin-labeled mouse anti-HM1.24 antibody was measured by the Cell-ELISA using WISH cells. Cell-ELISA plates were prepared as described above. After blocking, 50  $\mu$ l of

serial dilutions of the natural humanized anti-HM1.24 antibody that was obtained from the concentrate of the culture supernatant of COS-7 cells was added to each well, and 50  $\mu$ l of 2  $\mu$ g/ml biotin-labeled mouse anti-HM1.24 antibody was added simultaneously. After incubating at room temperature for two hours and washing, peroxidase-labeled streptoavidin (manufactured by DAKO) was added. After incubating at room temperature for one hour and washing, the reaction was stopped by adding 50  $\mu$ l of 6N sulfuric acid, and absorbance at 490 nm was measured using the MICROPLATE READER Model 3550 (manufactured by Bio-Rad).

(2) Antigen binding activity and binding inhibition activity

The evaluation of the H chain of natural humanized anti-HM1.24 antibody was conducted by measurement of the above-mentioned antigen binding activity and binding inhibition activity in combination with the L chain version a. The result, as shown in Figure 29 and 30, indicated that natural humanized anti-HM1.24 antibody (the secondary design antibody) has antigen binding activity and binding inhibition activity of a similar degree to the primary design antibody (reshaped human anti-HM1.24 antibody: the H chain version r).

(3) Measurement of the ADCC activity

ADCC (Antibody-dependent Cellular Cytotoxicity) activity was measured according to the method described in Reference Example 8.

1. Preparation of effector cells

To the peripheral blood of healthy human subject was added an equal amount of PBS(-), onto which Ficoll-Paque (manufactured by Pharmacia) was layered, and was centrifuged at 500 g for 30 minutes. The monocyte layer was taken therefrom and was washed twice with RPMI 1640 (manufactured by GIBCO BRL) supplemented with 10%

bovine fetal serum (manufactured by GIBCO BRL), and was adjusted to a cell density of  $5 \times 10^6$ /ml with the same culture liquid.

## 2. Preparation of target cells

5           The human myeloma cell line KPMM2 (Deposit No. P-14170, Patent application No. 6-58082) was radiolabeled by incubating in RPMI 1640 (manufactured by GIBCO BRL) supplemented with 10% bovine fetal serum (manufactured by GIBCO BRL) together with 0.1 mCi of  $^{51}\text{Cr}$ -sodium chromate  
10       at 37 °C for 60 minutes. After radiolabeling, cells were washed three times with the same buffer and adjusted to a concentration of  $2 \times 10^5$ /ml.

## 3. Measurement of ADCC assay

15           Into a 96-well U-bottomed plate (manufactured by Corning) were added 50  $\mu\text{l}$  of  $2 \times 10^5$  target cells/ml, 50  $\mu\text{l}$  of the antibody solution previously prepared at 4  $\mu\text{g}/\text{ml}$ , 0.4  $\mu\text{g}/\text{ml}$ , 0.04  $\mu\text{g}/\text{ml}$ , and 0.004  $\mu\text{g}/\text{ml}$ , and reacted at 4 °C for 15 minutes. A solution that does not contain natural humanized anti-HM1.24 antibody (the  
20       secondary design antibody) was similarly prepared and used as a control.

          Then, 100  $\mu\text{l}$  of  $5 \times 10^5$  effector cells/ml was added thereto, and cultured in a  $\text{CO}_2$ -incubator for 4 hours, wherein the ratio (E:T) of the effector cells (E)  
25       to the target cells (T) was set at 0:1, 20:1, and 50:1. Since the final concentration of each antibody was diluted by four-fold, they were 1  $\mu\text{g}/\text{ml}$ , 0.1  $\mu\text{g}/\text{ml}$ , 0.01  $\mu\text{g}/\text{ml}$ , and 0.001  $\mu\text{g}/\text{ml}$  as well as no antibody addition control.

30           One hundred  $\mu\text{l}$  of the supernatant was taken and the radioactivity released into the culture supernatant was measured by a gamma counter (ARC361, manufactured by Aloka). For measurement of the maximum radioactivity, 1% NP-40 (manufactured by Nacalai Tesque Inc.) was used.  
35       Cytotoxicity (%) was calculated by  $(A-C)/(B-C) \times 100$ ,

wherein A is radioactivity (cpm) released in the presence of antibody, B is radioactivity (cpm) released by NP-40, and C is radioactivity (cpm) released by the culture medium alone without antibody.

5           4. Result

As shown in Fig. 33, when the natural humanized anti-HM1.24 antibody (the secondary design antibody) was added, specific chromium release rate increased with the increase in the E:T ratio depending on antibody  
10 concentration as compared to the no antibody added control. This, therefore, indicated that this natural humanized anti-HM1.24 antibody (the secondary design antibody) has ADCC activity.

The present invention relates to a method of  
15 preparing natural humanized antibody and the natural humanized antibody obtained by said method of preparation. This is a highly excellent humanization technology that has solved the problems associated with CDR-grafting (Jones, P. T. et al., Nature (1986) 321,  
20 522-525) created by G. Winter. Construction of the primary design antibody may be considered as an intermediate stage for the construction of humanized antibody comprising natural human FRs. When antibody is developed as a pharmaceutical product comprising  
25 recombinant protein, natural humanized antibody that comprises naturally occurring human FRs is more excellent in terms of antigenicity and safety.

Effects of the Invention

30           Since the natural humanized antibody obtained by the method of preparation of the present invention does not contain the amino acid residues of non-naturally occurring artificial FRs that are contained in the humanized antibody produced by the conventional  
35 humanization technology, it is expected to have low antigenicity. Furthermore, it was shown that the natural humanized antibody obtained by the method of preparation

of the present invention has an activity similar to that  
of antibody derived from a non-human mammal that was used  
as a template for humanization. Therefore, the natural  
humanized antibody obtained by the method of preparation  
5 of the present invention is useful for therapeutic  
administration to humans.

Reference to the microorganisms deposited under the  
Patent Cooperation Treaty, Rule 13-2, and the name of the  
10 Depository Institute

Depository Institute

Name: the National Institute of Bioscience and Human  
Technology, Agency of Industrial Science and Technology

Address: 1-3, Higashi 1-chome, Tsukuba-shi, Ibaraki,  
15 Japan

Organism (1)

Indication: Escherichia coli DH5 $\alpha$  (pRS38-pUC19)

Accession number: FERM BP-4434

Deposition Date: October 5, 1993

20 Organism (2)

Indication: Hybridoma HM1.24

Accession number: FERM BP-5233

Deposition Date: September 14, 1995

Organism (3)

25 Indication: Escherichia coli DH5 $\alpha$  (pUC19-RVHr-AHM-gyl)

Accession number: FERM BP-5643

Deposition Date: August 29, 1996

Organism (4)

Indication: Escherichia coli DH5 $\alpha$  (pUC19-1.24H-gyl)

30 Accession number: FERM BP-5644

Deposition Date: August 29, 1996

Organism (5)

Indication: Escherichia coli DH5 $\alpha$  (pUC19-RVLa-AHM-gk)

Accession number: FERM BP-5645

35 Deposition Date: August 29, 1996

Organism (6)





# SEQUENCE LISTING

Sequence: 1

Sequence length: 394

Sequence type: Nucleic acid

Topology: Linear

Molecular type: cDNA

Sequence:

ATG GGC TTC AAG ATG GAG TCA CAT TTT CTG GTC TTT GTA TTC GTG TTT	48
Met Gly Phe Lys Met Glu Ser His Phe Leu Val Phe Val Phe Val Phe	
-20 -15 -10	
CTC TGG TTG TCT GGT GTT GAC GGA GAC ATT GTG ATG ACC CAG TCT CAC	96
Leu Trp Leu Ser Gly Val Asp Gly Asp Ile Val Met Thr Gln Ser His	
-5 -1 1 5	
AAA TTC ATG TCC ACA TCA GTA GGA GAC AGG GTC AGC ATC ACC TGC AAG	144
Lys Phe Met Ser Thr Ser Val Gly Asp Arg Val Ser Ile Thr Cys Lys	
10 15 20	
GCC AGT CAG GAT GTG AAT ACT GCT GTA GCC TGG TAT CAA CAA AAA CCA	192
Ala Ser Gln Asp Val Asn Thr Ala Val Ala Trp Tyr Gln Gln Lys Pro	
25 30 35 40	
GGA CAA TCG CCT AAA CTA CTG ATT TAC TCG GCA TCC AAC CGG TAC ACT	240
Gly Gln Ser Pro Lys Leu Leu Ile Tyr Ser Ala Ser Asn Arg Tyr Thr	
45 50 55	
GGA GTC CCT GAT CGC ATC ACT GGC AGT GGA TCT GGG ACG GAT TTC ACT	288
Gly Val Pro Asp Arg Ile Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr	
60 65 70	
TTC ACC ATC AGC AGT GTG CAG GCG GAA GAC CTG GCA CTT TAT TAC TGT	336
Phe Thr Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Leu Tyr Tyr Cys	
75 80 85	
CAG CAA CAT TAT AGT ACT CCA TTC ACG TTC GGC TCG GGG ACA AAG TTG	384
Gln Gln His Tyr Ser Thr Pro Phe Thr Phe Gly Ser Gly Thr Lys Leu	
90 95 100	
GAA ATA AAA C	394
Glu Ile Lys	
105	

Sequence: 3

Sequence length: 418

002220 00000000



Lys Ala Ser Gln Asp Val Asn Thr Ala Val Ala  
1 5 10

Sequence: 6

Sequence length: 7

Sequence type: Amino acid

Topology: Linear

Molecular type: Peptide

Sequence:

Ser Ala Ser Asn Arg Tyr Thr  
1 5

Sequence: 7

Sequence length: 9

Sequence type: Amino acid

Topology: Linear

Molecular type: Peptide

Sequence:

Gln Gln His Tyr Ser Thr Pro Phe Thr  
1 5

Sequence: 8

Sequence length: 5

Sequence type: Amino acid

Topology: Linear

Molecular type: Peptide

Sequence:

Pro Tyr Trp Met Gln  
1 5

Sequence: 9

Sequence length: 16

Sequence type: Amino acid

Topology: Linear

Molecular type: Peptide

Sequence:

Ser Ile Phe Gly Asp Gly Asp Thr Arg Tyr Ser Gln Lys Phe Lys Gly  
1 5 10 15

Sequence: 10

Sequence length: 11

Sequence type: Amino acid

Topology: Linear

Molecular type: Peptide

Sequence:

Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr

1 5 10

Sequence: 11

Sequence length: 379

Sequence type: Nucleic acid

Topology: Linear

Molecular type: cDNA

Sequence:

ATG GGA TGG AGC TGT ATC ATC CTC TCC TTG GTA GCA ACA GCT ACA GGT	48
Met Gly Trp Ser Cys Ile Ile Leu Ser Leu Val Ala Thr Ala Thr Gly	
-15 -10 -5	
GTC CAC TCC GAC ATC CAG ATG ACC CAG AGC CCA AGC AGC CTG AGC GCC	96
Val His Ser Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala	
-1 1 5 10	
AGC GTG GGT GAC AGA GTG ACC ATC ACC TGT AAG GCT AGT CAG GAT GTG	144
Ser Val Gly Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Val	
15 20 25	
AAT ACT GCT GTA GCC TGG TAC CAG CAG AAG CCA GGA AAG GCT CCA AAG	192
Asn Thr Ala Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys	
30 35 40 45	
CTG CTG ATC TAC TCG GCA TCC AAC CGG TAC ACT GGT GTG CCA AGC AGA	240
Leu Leu Ile Tyr Ser Ala Ser Asn Arg Tyr Thr Gly Val Pro Ser Arg	
50 55 60	
TTC AGC GGT AGC GGT AGC GGT ACC GAC TTC ACC TTC ACC ATC AGC AGC	288
Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser	
65 70 75	
CTC CAG CCA GAG GAC ATC GCT ACC TAC TAC TGC CAG CAA CAT TAT AGT	336
Leu Gln Pro Glu Asp Ile Ala Thr Tyr Tyr Cys Gln Gln His Tyr Ser	
80 85 90	
ACT CCA TTC ACG TTC GGC CAA GGG ACC AAG GTG GAA ATC AAA C	379
Thr Pro Phe Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys	
95 100 105	

Sequence: 13

0022250 35550550

Sequence length: 379

Sequence type: Nucleic acid

Topology: Linear

Molecular type: cDNA

Sequence:

ATG GGA TGG AGC TGT ATC ATC CTC TCC TTG GTA GCA ACA GCT ACA GGT	48
Met Gly Trp Ser Cys Ile Ile Leu Ser Leu Val Ala Thr Ala Thr Gly	
-15 -10 -5	
GTC CAC TCC GAC ATC CAG ATG ACC CAG AGC CCA AGC AGC CTG AGC GCC	96
Val His Ser Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala	
-1 1 5 10	
AGC GTG GGT GAC AGA GTG ACC ATC ACC TGT AAG GCT AGT CAG GAT GTG	144
Ser Val Gly Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Val	
15 20 25	
AAT ACT GCT GTA GCC TGG TAC CAG CAG AAG CCA GGA AAG GCT CCA AAG	192
Asn Thr Ala Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys	
30 35 40 45	
CTG CTG ATC TAC TCG GCA TCC AAC CGG TAC ACT GGT GTG CCA AGC AGA	240
Leu Leu Ile Tyr Ser Ala Ser Asn Arg Tyr Thr Gly Val Pro Ser Arg	
50 55 60	
TTC AGC GGT AGC GGT AGT GGT ACC GAC TAC ACC TTC ACC ATC AGC AGC	288
Phe Ser Gly Ser Gly Ser Gly Thr Asp Tyr Thr Phe Thr Ile Ser Ser	
65 70 75	
CTC CAG CCA GAG GAC ATC GCT ACC TAC TAC TGC CAG CAA CAT TAT AGT	336
Leu Gln Pro Glu Asp Ile Ala Thr Tyr Tyr Cys Gln Gln His Tyr Ser	
80 85 90	
ACT CCA TTC ACG TTC GGC CAA GGG ACC AAG GTG GAA ATC AAA C	379
Thr Pro Phe Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys	
95 100 105	

Sequence: 15

Sequence length: 418

Sequence type: Nucleic acid

Topology: Linear

Molecular type: cDNA

Sequence:

ATG GAC TGG ACC TGG AGG GTC TTC TTC TTG CTG GCT GTA GCT CCA GGT	48
Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly	
-15 -10 -5	
GCT CAC TCC CAG GTG CAG CTG GTG CAG TCT GGG GCT GAG GTG AAG AAG	96
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys	
-1 1 5 10	
CCT GGG GCC TCA GTG AAG GTT TCC TGC AAG GCA TCT GGA TAC ACC TTC	144
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe	
15 20 25	
ACT CCC TAC TGG ATG CAG TGG GTG CGA CAG GCC CCT GGA CAA GGG CTT	192
Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu	
30 35 40 45	
GAG TGG ATG GGA TCT ATT TTT CCT GGA GAT GGT GAT ACT AGG TAC AGT	240
Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser	
50 55 60	
CAG AAG TTC AAG GGC AGA GTC ACC ATG ACC GCA GAC ACG TCC ACG AGC	288
Gln Lys Phe Lys Gly Arg Val Thr Met Thr Ala Asp Thr Ser Thr Ser	
65 70 75	
ACA GTC TAC ATG GAG CTG AGC AGC CTG AGA TCT GAG GAC ACG GCC GTG	336
Thr Val Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val	
80 85 90	
TAT TAC TGT GCG AGA GGA TTA CGA CGA GGG GGG TAC TAC TTT GAC TAC	384
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr	
95 100 105	
TGG GGG CAA GGG ACC ACG GTC ACC GTC TCC TCA G	418
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser	
110 115 120	

Sequence: 17

Sequence length: 418

Sequence type: Nucleic acid

Topology: Linear

Molecular type: cDNA

Sequence:

ATG GAC TGG ACC TGG AGG GTC TTC TTC TTG CTG GCT GTA GCT CCA GGT	48
Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly	
-15 -10 -5	

GCT CAC TCC CAG GTG CAG CTG GTG CAG TCT GGG GCT GAG GTG AAG AAG	96
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys	
-1 1 5 10	
CCT GGG GCC TCA GTG AAG GTT TCC TGC AAG GCA TCT GGA TAC ACC TTC	144
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe	
15 20 25	
ACT CCC TAC TGG ATG CAG TGG GTG CGA CAG GCC CCT GGA CAA GGG CTT	192
Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu	
30 35 40 45	
GAG TGG ATG GGA TCT ATT TTT CCT GGA GAT GGT GAT ACT AGG TAC AGT	240
Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser	
50 55 60	
CAG AAG TTC AAG GGC AAA GTC ACC ATG ACC GCA GAC ACG TCC ACG AGC	288
Gln Lys Phe Lys Gly Lys Val Thr Met Thr Ala Asp Thr Ser Thr Ser	
65 70 75	
ACA GTC TAC ATG GAG CTG AGC AGC CTG AGA TCT GAG GAC ACG GCC GTG	336
Thr Val Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val	
80 85 90	
TAT TAC TGT GCG AGA GGA TTA CGA CGA GGG GGG TAC TAC TTT GAC TAC	384
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr	
95 100 105	
TGG GGG CAA GGG ACC ACG GTC ACC GTC TCC TCA G	418
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser	
110 115 120	

Sequence: 19

Sequence length: 418

Sequence type: Nucleic acid

Topology: Linear

Molecular type: cDNA

Sequence:

ATG GAC TGG ACC TGG AGG GTC TTC TTC TTG CTG GCT GTA GCT CCA GGT	48
Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly	
-15 -10 -5	
GCT CAC TCC CAG GTG CAG CTG GTG CAG TCT GGG GCT GAG GTG AAG AAG	96
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys	
-1 1 5 10	

002250-35060560









CAG AAG TTC AAG GGC AGA GCC ACC CTG ACT GCA GAC ACG TCC TCG AGC	288
Gln Lys Phe Lys Gly Arg Ala Thr Leu Thr Ala Asp Thr Ser Ser Ser	
65 70 75	
ACA GCC TAC ATG GAG CTG AGC AGC CTG AGA TCT GAG GAC ACG GCC GTG	336
Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val	
80 85 90	
TAT TAC TGT GCG AGA GGA TTA CGA CGA GGG GGG TAC TAC TTT GAC TAC	384
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr	
95 100 105	
TGG GGG CAA GGG ACC ACG GTC ACC GTC TCC TCA G	418
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser	
110 115 120	

Sequence: 27

Sequence length: 418

Sequence type: Nucleic acid

Topology: Linear

Molecular type: cDNA

Sequence:

ATG GAC TGG ACC TGG AGG GTC TTC TTC TTG CTG GCT GTA GCT CCA GGT	48
Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly	
-15 -10 -5	
GCT CAC TCC CAG GTG CAG CTG GTG CAG TCT GGG GCT GAG GTG AAG AAG	96
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys	
-1 1 5 10	
CCT GGG GCC TCA GTG AAG GTT TCC TGC AAG GCA TCT GGA TAC ACC TTC	144
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe	
15 20 25	
ACT CCC TAC TGG ATG CAG TGG GTG CGA CAG CGC CCT GGA CAA GGG CTT	192
Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Arg Pro Gly Gln Gly Leu	
30 35 40 45	
GAG TGG ATG GGA TCT ATT TTT CCT GGA GAT GGT GAT ACT AGG TAC AGT	240
Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser	
50 55 60	
CAG AAG TTC AAG GGC AGA GTC ACC ATG ACC GCA GAC ACG TCC ACG AGC	288
Gln Lys Phe Lys Gly Arg Val Thr Met Thr Ala Asp Thr Ser Thr Ser	
65 70 75	

002250 55050550







Sequence type: Nucleic acid

Topology: Linear

Molecular type: cDNA

Sequence:

ATG GAC TGG ACC TGG AGG GTC TTC TTC TTG CTG GCT GTA GCT CCA GGT	48
Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly	
-15 -10 -5	
GCT CAC TCC CAG GTG CAG CTG GTG CAG TCT GGG GCT GAG GTG AAG AAG	96
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys	
-1 1 5 10	
CCT GGG GCC TCA GTG AAG GTT TCC TGC AAG GCA TCT GGA TAC ACC TTC	144
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe	
15 20 25	
ACT CCC TAC TGG ATG CAG TGG GTG CGA CAG GCC CCT GGA CAA GGG CTT	192
Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu	
30 35 40 45	
GAG TGG ATG GGA TCT ATT TTT CCT GGA GAT GGT GAT ACT AGG TAC AGT	240
Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser	
50 55 60	
CAG AAG TTC AAG GGC AAA GTC ACC ATG ACC GCA GAC ACG TCC TCG AGC	288
Gln Lys Phe Lys Gly Lys Val Thr Met Thr Ala Asp Thr Ser Ser Ser	
65 70 75	
ACA GCC TAC ATG CAG CTG AGC AGC CTA AGA TCT GAG GAC ACG GCC GTG	336
Thr Ala Tyr Met Gln Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val	
80 85 90	
TAT TAC TGT GCG AGA GGA TTA CGA CGA GGG GGG TAC TAC TTT GAC TAC	384
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr	
95 100 105	
TGG GGG CAA GGG ACC ACG GTC ACC GTC TCC TCA G	418
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser	
110 115 120	

Sequence: 37

Sequence length: 418

Sequence type: Nucleic acid

Topology: Linear

Molecular type: cDNA

Sequence:

002250 86060560



ATG GAC TGG ACC TGG AGG GTC TTC TTC TTG CTG GCT GTA GCT CCA GGT	48
Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly	
-15 -10 -5	
GCT CAC TCC CAG GTG CAG CTG GTG CAG TCT GGG GCT GAG GTG AAG AAG	96
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys	
-1 1 5 10	
CCT GGG GCC TCA GTG AAG GTT TCC TGC AAG GCA TCT GGA TAC ACC TTC	144
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe	
15 20 25	
ACT CCC TAC TGG ATG CAG TGG GTG CGA CAG GCC CCT GGA CAA GGG CTT	192
Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu	
30 35 40 45	
GAG TGG ATG GGA TCT ATT TTT CCT GGA GAT GGT GAT ACT AGG TAC AGT	240
Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser	
50 55 60	
CAG AAG TTC AAG GGC AAA GTC ACC ATG ACC GCA GAC ACG TCC TCG AGC	288
Gln Lys Phe Lys Gly Lys Val Thr Met Thr Ala Asp Thr Ser Ser Ser	
65 70 75	
ACA GCC TAC ATG CAG CTG AGC ATC CTG AGA TCT GAG GAC ACG GCC GTG	336
Thr Ala Tyr Met Gln Leu Ser Ile Leu Arg Ser Glu Asp Thr Ala Val	
80 85 90	
TAT TAC TGT GCG AGA GGA TTA CGA CGA GGG GGG TAC TAC TTT GAC TAC	384
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr	
95 100 105	
TGG GGG CAA GGG ACC ACG GTC ACC GTC TCC TCA G	418
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser	
110 115 120	

Sequence: 39

Sequence length: 418

Sequence type: Nucleic acid

Topology: Linear

Molecular type: cDNA

Sequence:

ATG GAC TGG ACC TGG AGG GTC TTC TTC TTG CTG GCT GTA GCT CCA GGT	48
Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly	
-15 -10 -5	

GCT CAC TCC CAG GTG CAG CTG GTG CAG TCT GGG GCT GAG GTG AAG AAG	96
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys	
-1 1 5 10	
CCT GGG GCC TCA GTG AAG GTT TCC TGC AAG GCA TCT GGA TAC ACC TTC	144
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe	
15 20 25	
ACT CCC TAC TGG ATG CAG TGG GTG CGA CAG GCC CCT GGA CAA GGG CTT	192
Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu	
30 35 40 45	
GAG TGG ATG GGA TCT ATT TTT CCT GGA GAT GGT GAT ACT AGG TAC AGT	240
Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser	
50 55 60	
CAG AAG TTC AAG GGC AAA GTC ACC ATG ACC GCA GAC ACG TCC TCG AGC	288
Gln Lys Phe Lys Gly Lys Val Thr Met Thr Ala Asp Thr Ser Ser Ser	
65 70 75	
ACA GCC TAC ATG CAG CTG AGC ATC CTG AGA TCT GAG GAC TCG GCC GTG	336
Thr Ala Tyr Met Gln Leu Ser Ile Leu Arg Ser Glu Asp Ser Ala Val	
80 85 90	
TAT TAC TGT GCG AGA GGA TTA CGA CGA GGG GGG TAC TAC TTT GAC TAC	384
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr	
95 100 105	
TGG GGG CAA GGG ACC ACG GTC ACC GTC TCC TCA G	418
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser	
110 115 120	

Sequence: 41

Sequence length: 418

Sequence type: Nucleic acid

Topology: Linear

Molecular type: cDNA

Sequence:

ATG GAC TGG ACC TGG AGG GTC TTC TTC TTG CTG GCT GTA GCT CCA GGT	48
Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly	
-15 -10 -5	
GCT CAC TCC CAG GTG CAG CTG GTG CAG TCT GGG GCT GAG GTG AAG AAG	96
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys	
-1 1 5 10	







CAG AAG TTC AAG GGC AGA GTC ACC ATG ACC GCA GAC ACG TCC TCG AGC	288
Gln Lys Phe Lys Gly Arg Val Thr Met Thr Ala Asp Thr Ser Ser Ser	
65 70 75	
ACA GTC TAC ATG GAG CTG AGC AGC CTG AGA TCT GAG GAC ACG GCC GTG	336
Thr Val Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val	
80 85 90	
TAT TAC TGT GCG AGA GGA TTA CGA CGA GGG GGG TAC TAC TTT GAC TAC	384
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr	
95 100 105	
TGG GGG CAA GGG ACC ACG GTC ACC GTC TCC TCA G	418
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser	
110 115 120	

Sequence: 49

Sequence length: 418

Sequence type: Nucleic acid

Topology: Linear

Molecular type: cDNA

Sequence:

ATG GAC TGG ACC TGG AGG GTC TTC TTC TTG CTG GCT GTA GCT CCA GGT	48
Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly	
-15 -10 -5	
GCT CAC TCC CAG GTG CAG CTG GTG CAG TCT GGG GCT GAG GTG AAG AAG	96
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys	
-1 1 5 10	
CCT GGG GCC TCA GTG AAG GTT TCC TGC AAG GCA TCT GGA TAC ACC TTC	144
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe	
15 20 25	
ACT CCC TAC TGG ATG CAG TGG GTG CGA CAG GCC CCT GGA CAA GGG CTT	192
Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu	
30 35 40 45	
GAG TGG ATG GGA TCT ATT TTT CCT GGA GAT GGT GAT ACT AGG TAC AGT	240
Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser	
50 55 60	
CAG AAG TTC AAG GGC AGA GTC ACC ATG ACC GCA GAC AAG TCC ACG AGC	288
Gln Lys Phe Lys Gly Arg Val Thr Met Thr Ala Asp Lys Ser Thr Ser	
65 70 75	

ACA GCC TAC ATG GAG CTG AGC AGC CTG AGA TCT GAG GAC ACG GCC GTG 336  
 Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val  
 80 85 90

TAT TAC TGT GCG AGA GGA TTA CGA CGA GGG GGG TAC TAC TTT GAC TAC 384  
 Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
 95 100 105

TGG GGG CAA GGG ACC ACG GTC ACC GTC TCC TCA G 418  
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 110 115 120

Sequence: 51

Sequence length: 40

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

ACTAGTCGAC ATGAAGTTGC CTGTTAGGCT GTTGGTGCTG 40

Sequence: 52

Sequence length: 39

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

ACTAGTCGAC ATGGAGWCAG ACACACTCCT GYTATGGGT 39

Sequence: 53

Sequence length: 40

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

ACTAGTCGAC ATGAGTGTGC TCACTCAGGT CCTGGS GTTG 40

Sequence: 54

Sequence length: 43

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

ACTAGTCGAC ATGAGGRCCC CTGCTCAGWT TYTTGGMWTC TTG 43

002220-25050550





Sequence length: 37

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

ACTAGTCGAC ATGTATATAT GTTTGTTGTC TATTTCT

37

Sequence: 61

Sequence length: 38

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

ACTAGTCGAC ATGGAAGCCC CAGCTCAGCT TCTCTTCC

38

Sequence: 62

Sequence length: 27

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GGATCCCGGG TGGATGGTGG GAAGATG

27

Sequence: 63

Sequence length: 25

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

TAGAGTCACC GAGGAGCCAG TTGTA

25

Sequence: 64

Sequence length: 26

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GGATCCCGGG AGTGGATAGA CCGATG

26

Sequence: 65

Sequence length: 34

002220" 86060560

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GATAAGCTTC CACCATGGGC TTCAAGATGG AGTC

34

Sequence: 66

Sequence length: 34

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GATAAGCTTC CACCATGGAA TGTAAGTGA TACT

34

Sequence: 67

Sequence length: 34

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GGCGGATCCA CTCACGTTTT ATTTCCAAGT TTGT

34

Sequence: 68

Sequence length: 34

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GGCGGATCCA CTCACCTGAG GAGACTGTGA GAGT

34

Sequence: 69

Sequence length: 18

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

CAGACAGTGG TTCAAAGT

18

Sequence: 70

Sequence length: 26

Sequence type: Nucleic acid

002250"85050350

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GAATTCGGAT CCACTCACGT TTGATT

26

Sequence: 71

Sequence length: 48

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

AGTCAGGATG TGAATACTGC TGTAGCCTGG TACCAGCAGA AGCCAGGA

48

Sequence: 72

Sequence length: 39

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GCATCCAACC GGTACACTGG TGTGCCAAGC AGATTCAGC

39

Sequence: 73

Sequence length: 45

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

CAACATTATA GTACTCCATT CACGTTTCGGC CAAGGGACCA AGGTG

45

Sequence: 74

Sequence length: 47

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GCAGTATTCA CATCCTGACT GGCCTTACAG GTGATGGTCA CTCTGTC

47

Sequence: 75

Sequence length: 38

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

ACACCAAGTGT ACCGGTTGGA TGCCGAGTAG ATCAGCAG

38

Sequence: 76

Sequence length: 41

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GTGAATGGAG TACTATAATG TTGCTGGCAG TAGTAGGTAG C

41

Sequence: 77

Sequence length: 31

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GGTACCGACT ACACCTTCAC CATCAGCAGC C

31

Sequence: 78

Sequence length: 31

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GGTGAAGGTG TAGTCGGTAC CGCTACCGCT A

31

Sequence: 79

Sequence length: 144

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

ATGCCTTGCA GGAAACCTTC ACTGAGGCC CAGGCTTCTT CACCTCAGCC CCAGACTGCA

60

CCAGCTGCAC CTGGGAGTGA GCACCTGGAG CTACAGCCAG CAAGAAGAAG ACCCTCCAGG

120

TCCAGTCCAT GGTGGAAGCT TATC

144

Sequence: 80

Sequence length: 130

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

TCAGTGAAGG TTCCTGCAA GGCATCTGGA TACACCTTCA CTCCCTACTG GATGCAGTGG	60
GTGCGACAGG CCCCTGGACA AGGGCTTGAG TGGATGGGAT CTATTTTTC TGGAGATGGT	120
GATACTAGGT	130

Sequence: 81

Sequence length: 131

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

AATACACGGC CGTGTCTCA GATCTCAGGC TGCTCAGCTC CATGTAGACT GTGCTCGTGG	60
ACGTGTCTGC GGTTCATGGTG ACTCTGCCCT TGAACCTCTG ACTGTACCTA GTATCACCAT	120
CTCCAGGAAA A	131

Sequence: 82

Sequence length: 119

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GAGATCTGAG GACACGGCCG TGTATTACTG TGCGAGAGGA TTACGACGAG GGGGGTACTA	60
CTTTGACTAC TGGGGGCAAG GGACCACGGT CACCGTCTCC TCAGGTGAGT GGATCCGAC	119

Sequence: 83

Sequence length: 25

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GATAAGCTTC CACCATGGAC TGGAC	25
-----------------------------	----

Sequence: 84

Sequence length: 25

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GTCGGATCCA CTCACCTGAG GAGAC

25

Sequence: 85

Sequence length: 26

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

AAGTTCAAGG GCAAAGTCAC CATGAC

26

Sequence: 86

Sequence length: 26

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GTCATGGTGA CTTTGCCCTT GAACTT

26

Sequence: 87

Sequence length: 26

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

ATGACCGCAG ACAAGTCCAC GAGCAC

26

Sequence: 88

Sequence length: 26

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GTGCTCGTGG ACTTGTCTGC GGTCAT

26

Sequence: 89

Sequence length: 47

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

AAGTTCAAGG GCAAAGTCAC CATGACCGCA GACAAGTCCA CGAGCAC

47

Sequence: 90

Sequence length: 47

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GTGCTCGTGG ACTTGCTCTGC GGTCAATGGTG ACTTTGCCCT TGAAGTT

47

Sequence: 91

Sequence length: 38

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

AAGTTCAAGG GCAGAGCCAC CCTGACCGCA GACACGTC

38

Sequence: 92

Sequence length: 38

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GACGTGTCTG CGGTCAGGGT GGCTCTGCCC TTGAAGTT

38

Sequence: 93

Sequence length: 18

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

CAGACAGTGG TTCAAAGT

18

Sequence: 94

Sequence length: 17

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GCCCCAAGC CAAGGTC

17

Sequence: 95

Sequence length: 23  
Sequence type: Nucleic acid  
Topology: Linear  
Molecular type: Synthetic DNA  
Sequence:

ATTTTCCTG GAGATGGTGA TAC 23

Sequence: 96

Sequence length: 23  
Sequence type: Nucleic acid  
Topology: Linear  
Molecular type: Synthetic DNA  
Sequence:

GTATCACCAT CTCCAGGAAA TAT 23

Sequence: 97

Sequence length: 418  
Sequence type: Nucleic acid  
Topology: Linear  
Molecular type: cDNA  
Sequence:

ATG GAA TGT AAC TGG ATA CTT CCT TTT ATT CTG TCA GTA ACT TCA GGT 48

Met Glu Cys Asn Trp Ile Leu Pro Phe Ile Leu Ser Val Thr Ser Gly  
-15 -10 -5

GCC TAC TCA CAG GTT CAA CTC CAG CAG TCT GGG GCT GAG CTG GCA AGA 96

Ala Tyr Ser Gln Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg  
-1 1 5 10

CCT GGG GCT TCA GTG AAG TTG TCC TGC AAG GCT TCT GGC TAC ACC TTT 144

Pro Gly Ala Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
15 20 25

ACT CCC TAC TGG ATG CAG TGG GTA AAA CAG AGG CCT GGA CAG GGT CTG 192

Thr Pro Tyr Trp Met Gln Trp Val Lys Gln Arg Pro Gly Gln Gly Leu  
30 35 40 45

GAA TGG ATT GGG TCT ATT TTT CCT GGA GAT GGT GAT ACT AGG TAC AGT 240

Glu Trp Ile Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
50 55 60

CAG AAG TTC AAG GGC AGA GTC ACC ATG ACC GCA GAC ACG TCC ACG AGC 288

Gln Lys Phe Lys Gly Arg Val Thr Met Thr Ala Asp Thr Ser Thr Ser  
65 70 75



ACA GTC TAC ATG GAG CTG AGC AGC CTG AGA TCT GAG GAC ACG GCC GTG	336
Thr Val Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val	
80 85 90	
TAT TAC TGT GCG AGA GGA TTA CGA CGA GGG GGG TAC TAC TTT GAC TAC	384
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr	
95 100 105	
TGG GGG CAA GGG ACC ACG GTC ACC GTC TCC TCA G	418
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser	
110 115 120	

Sequence: 99

Sequence length: 418

Sequence type: Nucleic acid

Topology: Linear

Molecular type: cDNA

Sequence:

ATG GAC TGG ACC TGG AGG GTC TTC TTC TTG CTG GCT GTA GCT CCA GGT	48
Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly	
-15 -10 -5	
GCT CAC TCC CAG GTG CAG CTG GTG CAG TCT GGG GCT GAG GTG AAG AAG	96
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys	
-1 1 5 10	
CCT GGG GCC TCA GTG AAG GTT TCC TGC AAG GCA TCT GGA TAC ACC TTC	144
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe	
15 20 25	
ACT CCC TAC TGG ATG CAG TGG GTG CGA CAG GCC CCT GGA CAA GGG CTT	192
Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu	
30 35 40 45	
GAG TGG ATG GGA TCT ATT TTT CCT GGA GAT GGT GAT ACT AGG TAC AGT	240
Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser	
50 55 60	
CAG AAG TTC AAG GGC AAG GCC ACA TTG ACT GCA GAT AAA TCC TCC AGT	288
Gln Lys Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser	
65 70 75	
ACA GCC TAC ATG CAA CTC AGC ATC TTG GCA TTT GAG GAC TCT GCG GTC	336
Thr Ala Tyr Met Gln Leu Ser Ile Leu Ala Phe Glu Asp Ser Ala Val	
80 85 90	

TAT TAC TGT GCA AGA GGA TTA CGA CGA GGG GGG TAC TAC TTT GAC TAC 384  
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
95 100 105

TGG GGC CAA GGC ACC ACT CTC ACA GTC TCC TCA G 418  
Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser  
110 115 120

Sequence: 101

Sequence length: 38

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

CTGGTTCGGC CCACCTCTGA AGGTTCCAGA ATCGATAG 38

Sequence: 102

Sequence length: 35

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GCAGACACGT CCTCGAGCAC AGCCTACATG GAGCT 35

Sequence: 103

Sequence length: 35

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

AGCTCCATGT AGGCTGTGCT CGAGGACGTG TCTGC 35

Sequence: 104

Sequence length: 26

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

TGGGTGCGAC AGCGCCCTGG ACAAGG 26

Sequence: 105

Sequence length: 26

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

CCTTGTCCAG GGCGCTGTCG CACCCA

26

Sequence: 106

Sequence length: 41

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

TACATGGAGC TGAGCAGCCT GGCATTGAG GACACGGCCG T

41

Sequence: 107

Sequence length: 41

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

ACGGCCGTGT CCTCAAATGC CAGGCTGCTC AGCTCCATGT A

41

Sequence: 108

Sequence length: 26

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

AAGTTCAAGG GCAAAGCCAC CCTGAC

26

Sequence: 109

Sequence length: 26

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GTCAGGGTGG CTTTGCCCTT GAACTT

26

Sequence: 110

Sequence length: 23

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GCCTACATGC AGCTGAGCAG CCT

23

Sequence: 111

Sequence length: 23

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

AGGCTGCTCA GCTGCATGTA GGC

23

Sequence: 112

Sequence length: 38

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GCCTACATGC AGCTGAGCAT CCTGAGATCT GAGGACAC

38

Sequence: 113

Sequence length: 35

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GATCTCAGGA TGCTCAGCTG CATGTAGGCT GTGCT

35

Sequence: 114

Sequence length: 50

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GCCTACATGC AGCTGAGCAT CCTGAGATCT GAGGACTCGG CCGTGTATTA

50

Sequence: 115

Sequence length: 50

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

ACGGCCGAGT CCTCAGATCT CAGGATGCTC AGCTGCATGT AGGCTGTGCT

50

Sequence: 116

Sequence length: 20

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GAGCTGAGCA TCCTGAGATC

20

Sequence: 117

Sequence length: 26

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GATCTCAGGA TGCTCAGCTC CATGTA

26

Sequence: 118

Sequence length: 20

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

AGATCTGAGG ACTCGGCCGT

20

Sequence: 119

Sequence length: 20

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

ACGGCCGAGT CCTCAGATCT

20

Sequence: 120

Sequence length: 35

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GCAGACACGT CCACGAGCAC AGCCTACATG GAGCT

35

Sequence: 121

Sequence length: 35

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

AGCTCCATGT AGGCTGTGCT CGTGGACGTG TCTGC

35

Sequence: 122

Sequence length: 35

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GCAGACACGT CCTCGAGCAC AGTCTACATG GAGCT

35

Sequence: 123

Sequence length: 35

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

AGCTCCATGT AGACTGTGCT CGAGGACGTG TCTGC

35

Sequence: 124

Sequence length: 26

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

AGAGTCACCA TCACCGCAGA CAAGTC

26

Sequence: 125

Sequence length: 26

Sequence type: Nucleic acid

Topology: Linear

Molecular type: Synthetic DNA

Sequence:

GACTTGCTCG CGGTGATGGT GACTCT

26

Sequence: 126

Sequence length: 418

Sequence type: Nucleic acid

Topology: Linear

Molecular type: cDNA

Sequence:

ATG GAC TGG ACC TGG AGG GTC TTC TTC TTG CTG GCT GTA GCT CCA GGT	48
Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly	
-15 -10 -5	
GCT CAC TCC CAG GTG CAG CTG GTG CAG TCT GGG GCT GAG GTG AAG AAG	96
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys	
-1 1 5 10	
CCT GGG GCC TCA GTG AAG GTT TCC TGC AAG GCA TCT GGA TAC ACC TTC	144
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe	
15 20 25	
ACT CCC TAC TGG ATG CAG TGG GTG CGA CAG GCC CCT GGA CAA GGG CTT	192
Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu	
30 35 40 45	
GAG TGG ATG GGA TCT ATT TTT CCT GGA GAT GGT GAT ACT AGG TAC AGT	240
Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser	
50 55 60	
CAG AAG TTC AAG GGC AGA GTC ACC ATC ACC GCA GAC AAG TCC ACG AGC	288
Gln Lys Phe Lys Gly Arg Val Thr Ile Thr Ala Asp Lys Ser Thr Ser	
65 70 75	
ACA GCC TAC ATG GAG CTG AGC AGC CTG AGA TCT GAG GAC ACG GCC GTG	336
Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val	
80 85 90	
TAT TAC TGT GCG AGA GGA TTA CGA CGA GGG GGG TAC TAC TTT GAC TAC	384
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr	
95 100 105	
TGG GGG CAA GGG ACC ACG GTC ACC GTC TCC TCA G	418
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser	
110 115 120	

Sequence: 128

Sequence length: 1013

Sequence type: Nucleic acid

002220"86060560

Sequence:

100



ATT GTG CTG CTG GGC CTC AGC GCT CTG CTG CAG TGA GATCCCAGGA	575
Ile Val Leu Leu Gly Leu Ser Ala Leu Leu Gln ***	
170	175
180	
AGCTGGCACA TCTTGGAAGG TCCGTCCTGC TCGGCTTTTC GCTTGAACAT TCCCTTGATC	635
TCATCAGTTC TGAGCGGGTC ATGGGGCAAC ACGGTTAGCG GGGAGAGCAC GGGGTAGCCG	695
GAGAAGGGCC TCTGGAGCAG GTCTGGAGGG GCCATGGGGC AGTCCTGGGT CTGGGGACAC	755
AGTCGGGTTG ACCCAGGGCT GTCTCCCTCC AGAGCCTCCC TCCGGACAAT GAGTCCCCCC	815
TCTTGTCTCC CACCCTGAGA TTGGGCATGG GGTGCGGTGT GGGGGGCATG TGCTGCCTGT	875
TGTTATGGGT TTTTTTTGCG GGGGGGTTG CTTTTTCTG GGGTCTTTGA GCTCCAAAAA	935
AATAAACT TCCTTTGAGG GAGAGCACAC CTAAAAAAA AAAAAAAAAA AAAAAAAAAA	995
AAAATTCGGG CGGCCGCC	1013

002250-50000000

422 Rec'd PCT/PTO 22 MAR 2000

CLAIM

1. A method, for preparing a natural humanized antibody, which comprises conducting a homology search for the FR of a primary design antibody and selecting a natural human FR retaining the artificial amino acid residues contained in the FR of the primary design antibody and having a homology therewith.

2. The method of preparing a natural humanized antibody according to claim 1, which comprises conducting a homology search for the FR of a primary design antibody, selecting a natural human FR retaining the artificial amino acid residues contained in the FR of the primary design antibody and having a homology therewith, and replacing one or a plurality of different amino acid residues between the FR of the primary design antibody and the selected natural human FR.

3. The method of preparation according to claim 1 or 2, wherein the primary design antibody comprises CDRs derived from a first animal species, and FRs derived from a second animal species and having artificial amino acid residues.

4. The method of preparation according to claim 3, wherein the first animal species is rat and the second animal species is human.

5. The method of preparation according to any of claims 1 to 4, wherein the artificial amino acid residues are derived from the FR of non-human antibody.

6. A natural humanized antibody obtained by a method of preparation according to any of the claims 1 to 5.

7. A natural humanized antibody containing CDR derived from a first animal species and the FR derived from a second animal species, characterized in that said FR comprises an amino acid sequence having amino acid residue different from the FR used for CDR-grafting by one or a plurality of amino acid residues and that said FR has been replaced with an FR derived from a second

422 Rec'd PCT/PTO 22 MAR 2000

8. The natural humanized antibody according to claim 7, wherein the first animal species is rat and the second animal species is human.

10. An expression vector comprising the DNA according to claim 9.

12. A method of preparing a natural humanized antibody, which comprises culturing the cells into which an expression vector comprising the DNA according to claim 9 has been introduced and recovering the desired natural humanized antibody from the culture of said cells.

13. A pharmaceutical composition comprising a natural humanized antibody.

[illegible]

ABSTRACT

A reshaped human anti-HM1.24 antibody comprising:

5 (A) L chains each comprising (1) a constant region  
of a human L chain, and (2) FRs of a human L chain, and  
CDRs of L chain of mouse anti-HM1.24 monoclonal antibody;  
and

10 (B) H chains each comprising (2) a constant region  
of a human H chain, and (2) FRs of a human H chain, and  
CDRs of H chain of mouse anti-HM1.24 monoclonal antibody.  
Since the majority of the reshaped human antibody is  
derived from human antibody and the CDR has a low  
antigenicity, the reshaped human antibody of the present  
invention has low antigenicity and therefore is very  
15 promising in medical and therapeutic applications.

002220 36060550

Fig.1




Fig.2




Fig.3




Fig.4






Fig.5






Fig.6

V REGION OF THE MOUSE ANTI-HM 1.24 ANTIBODY



V REGION OF THE H CHAIN OF THE RESHAPED HUMAN ANTI-HM 1.24 ANTIBODY VERSION "a"




FIRST PCR

(a-HYA)

(HYS-h)

ASSEMBLY

SECOND PCR



V REGION OF H CHAIN OF HUMAN MOUSE HYBRID






Fig.7

V REGION OF THE MOUSE ANTI-HM 1.24 ANTIBODY



V REGION OF THE H CHAIN OF THE RESHAPED HUMAN ANTI-HM 1.24 ANTIBODY VERSION "a"



FIRST PCR

(a-HYA)

(HYS-h)

ASSEMBLY

SECOND PCR

V REGION OF H CHAIN OF MOUSE HUMAN HYBRID




Fig.8




Fig.9




Fig.10




Fig.11




Fig.12






Fig.13




Fig.14




Fig.15




Fig.16




Fig.17




Fig.18




Fig.19




Fig.20






Fig.21




Fig.22




Fig.23




Fig.24




Fig.25




Fig.26




Fig.27




Fig.28








Fig.29



—□— RESHAPED HUMAN  
ANTI-HM1.24 ANTIBODY  
(PRIMARY DESIGN ANTIBODY)

---○--- NATURAL HUMANIZED ANTIBODY  
(SECONDARY DESIGN ANTIBODY)

Fig.30



—□— RESHAPED HUMAN  
ANTI-HM1.24 ANTIBODY  
(PRIMARY DESIGN ANTIBODY)  
---○--- NATURAL HUMANIZED ANTIBODY  
(SECONDARY DESIGN ANTIBODY)

Fig.31




Fig.32





Fig.33



- NATURAL HUMANIZED ANTI-HM1.24 ANTIBODY  
(SECONDARY DESIGN ANTIBODY) 1  $\mu\text{g/ml}$
- ▲— NATURAL HUMANIZED ANTI-HM1.24 ANTIBODY  
(SECONDARY DESIGN ANTIBODY) 0.1  $\mu\text{g/ml}$
- ◆— NATURAL HUMANIZED ANTI-HM1.24 ANTIBODY  
(SECONDARY DESIGN ANTIBODY) 0.01  $\mu\text{g/ml}$
- NATURAL HUMANIZED ANTI-HM1.24 ANTIBODY  
(SECONDARY DESIGN ANTIBODY) 0.001  $\mu\text{g/ml}$
- NO ANTIBODY ADDITION

## Declaration and Power of Attorney For Patent Application

## 特許出願宣言書及び委任状

## Japanese Language Declaration

## 日本語宣言書

下記の氏名の発明者として、私は以下の通り宣言します。 As a below named inventor, I hereby declare that:

私の住所、私書箱、国籍は下記の私の氏名の後に記載された通りです。

My residence, post office address and citizenship are as stated next to my name.

下記の名称の発明に関して請求範囲に記載され、特許出願している発明内容について、私が最初かつ唯一の発明者（下記の氏名が一つの場合）もしくは最初かつ共同発明者であると（下記の名称が複数の場合）信じています。

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

NATURAL HUMANIZED ANTIBODY

上記発明の明細書（下記の欄で x 印がついていない場合は、本書に添付）は、

the specification of which is attached hereto unless the following box is checked:

☐ 月 日に提出され、米国出願番号または特許協定条約国際出願番号を \_\_\_\_\_ とし、  
(該当する場合) \_\_\_\_\_ に訂正されました。

☐ was filed on October 2, 1998  
as United States Application Number or  
PCT International Application Number  
PCT/JP98/04469 and was amended on  
\_\_\_\_\_ (if applicable).

私は、特許請求範囲を含む上記訂正後の明細書を検討し、内容を理解していることをここに表明します。

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

私は、連邦規則法典第 37 編第 1 条 5 6 項に定義されるとおり、特許資格の有無について重要な情報を開示する義務があることを認めます。

I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56.

## Japanese Language Declaration (日本語宣言書)

私は、米国法典第35編119条(a)-(d)項又は365条(b)項に基づき下記の、米国以外の国の少なくとも一方国を指定している特許協力条約365(a)項に基づく国際出願、又は外国での特許出願もしくは発明者証の出願について外国優先権をここに主張するとともに、優先権を主張している、本出願の前に出願された特許または発明者証の外国出願を以下に、枠内をマークすることで、示しています。

I hereby claim foreign priority under Title 35, United States Code, Section 119(a)-(d) or 365(b) of any foreign application(s) for patent or inventor's certificate, or 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate, or PCT International application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)  
外国での先行出願

Priority Claimed  
優先権主張

9-271726 (Pat. Appln.) Japan  
(Number) (Country)  
(番号) (国名)

3/October/1997  
(Day/Month/Year Filed)  
(出願年月日)

☒ ☐  
Yes No  
はい いいえ

(Number) (Country)  
(番号) (国名)

(Day/Month/Year Filed)  
(出願年月日)

☐ ☐  
Yes No  
はい いいえ

私は、第35編米国法典第119条(e)項に基づいて下記の米  
国特許出願規定に記載された権利をここに主張いたします。

I hereby claim the benefit under Title 35, United States Code, Section 119(e) of any United States provisional application(s) listed below.

(Application No.) (Filing Date)  
(出願番号) (出願日)

(Application No.) (Filing Date)  
(出願番号) (出願日)

私は、下記の米国法典第35編120条に基づいて下記の米  
国特許出願に記載された権利、又は米国を指定している特許  
協力条約365条(c)に基づく権利をここに主張します。ま  
た、本出願の各請求範囲の内容が米国法典第35編112条  
第1項又は特許協力条約で規定された方法で先行する米国特  
許出願に開示されていない限り、その先行米国出願書提出日  
以降で本出願書の日本国内または特許協力条約国際提出日ま  
での期間中に入手された、連邦規則法典第37編1条56項  
で定義された特許資格の有無に関する重要な情報について開  
示義務があることを認識しています。

I hereby claim the benefit under Title 35, United States Code, Section 120 of any United States application(s), or 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of Title 35, United States Code Section 112, I acknowledge the duty to disclose information which is material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of application.

(Application No.) (Filing Date)  
(出願番号) (出願日)

(Status: Patented, Pending, Abandoned)  
(現況: 特許許可済、係属中、放棄済)

(Application No.) (Filing Date)  
(出願番号) (出願日)

(Status: Patented, Pending, Abandoned)  
(現況: 特許許可済、係属中、放棄済)

私は、私自身の知識に基づいて本宣言書中で私が行なう表  
明が真実であり、かつ私の入手した情報と私の信じるところ  
に基づく表明が全て真実であると信じていること、さらに故  
意になされた虚偽の表明及びそれと同等の行為は米国法典第  
18編第1001条に基づき、罰金または拘禁、もしくはそ  
の両方により処罰されること、そしてそのような故意による  
虚偽の声明を行なえば、出願した、又は既に許可された特許  
の有効性が失われることを認識し、よってここに上記のごと  
く宣言を致します。

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Japanese Language Declaration  
(日本語宣言書)

委任状: 私は下記の発明者として、本出願に関する一切の  
手続きを米特許商標局に対して遂行する弁理士または代理人  
として、下記の者を指名いたします。(弁護士、または代理  
人の氏名及び登録番号を明記のこと)

POWER OF ATTORNEY: As a named inventor, I hereby appoint  
the following attorney(s) and/or agent(s) to prosecute this  
application and transact all business in the Patent and Trademark  
Office connected therewith (list name and registration number).

Stephen A. Bent, Reg. No. 29,768  
David A. Blumenthal, Reg. No. 26,257  
William T. Ellis, Reg. No. 26,874  
John J. Feldhaus, Reg. No. 28,822  
Patricia D. Granados, Reg. No. 33,683  
John P. Isacson, Reg. No. 33,715  
Eugene M. Lee, Reg. No. 32,039  
Richard Linn, Reg. No. 25,144  
Peter G. Mack, Reg. No. 26,001

Brian J. McNamara, Reg. No. 32,789  
Sybil Meloy, Reg. No. 22,749  
George E. Quillin, Reg. No. 32,792  
Colin G. Sandercock, Reg. No. 31,298  
Bernhard D. Saxe, Reg. No. 28,665  
Charles F. Schill, Reg. No. 27,590  
Richard L. Schwaab, Reg. No. 25,479  
Arthur Schwartz, Reg. No. 22,115  
Harold C. Wegner, Reg. No. 25,258

## 書類送付先

Send Correspondence to:

Foley & Lardner  
3000 K Street, N.W.  
P.O. Box 25696  
Washington, DC 20007-8696

Foley & Lardner  
3000 K Street, N.W.  
P.O. Box 25696  
Washington, DC 20007-8696

直接電話連絡先: (名前及び電話番号)

Direct Telephone Calls to: (name and telephone number)

(202)672-5300

(202)672-5300

唯一または第一発明者名	1-00	Full name of sole or first inventor Masayuki Tsuchiya
発明者の署名	日付	Inventor's signature <i>Masayuki Tsuchiya</i> Date March 10, 2000
住所		Residence Gotenba-shi, Shizuoka, Japan JPX
国籍		Citizenship Japanese
私書箱		Post Office Address c/o CHUGAI SEIYAKU KABUSHIKI KATSHA, 135, Komakado 1-chome, Gotenba-shi, Shizuoka 412-8513, Japan
第二共同発明者		Full name of second joint inventor, if any
第二共同発明者の署名	日付	Second inventor's signature Date
住所		Residence
国籍		Citizenship
私書箱		Post Office Address

(第三以降の共同発明者についても同様に記載し、署名をす (Supply similar information and signature for third and  
ること) subsequent joint inventors.)



SEQUENCE LISTING

<100> Chugai Seiyaku Kabushiki Kaisha

<120> Natural Humanized Antibody

<130> F885/PCT

<150> JP 9-271726

<151> 1997-10-03

<160> 129

<210> 1

<211> 394

<212> DNA

<213> Mouse

<223> cDNA coding for L chain V region of anti-HM1.24 antibody

<400> 1

```

atg ggc ttc aag atg gag tca cat ttt ctg gtc ttt gta ttc gtg ttt      48
Met Gly Phe Lys Met Glu Ser His Phe Leu Val Phe Val Phe Val Phe
          -20                      -15                      -10

ctc tgg ttg tct ggt gtt gac gga gac att gtg atg acc cag tct cac      96
Leu Trp Leu Ser Gly Val Asp Gly Asp Ile Val Met Thr Gln Ser His
          -5                      -1    1                      5

aaa ttc atg tcc aca tca gta gga gac agg gtc agc atc acc tgc aag      144
Lys Phe Met Ser Thr Ser Val Gly Asp Arg Val Ser Ile Thr Cys Lys
          10                      15                      20

gcc agt cag gat gtg aat act gct gta gcc tgg tat caa caa aaa cca      192
Ala Ser Gln Asp Val Asn Thr Ala Val Ala Trp Tyr Gln Gln Lys Pro
          25                      30                      35                      40

gga caa tcg cct aaa cta ctg att tac tcg gca tcc aac cgg tac act      240
Gly Gln Ser Pro Lys Leu Leu Ile Tyr Ser Ala Ser Asn Arg Tyr Thr
          45                      50                      55

```

00222200 86060560

gga gtc cct gat cgc atc act ggc agt gga tct ggg acg gat ttc act 288  
 Gly Val Pro Asp Arg Ile Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr  
 60 65 70  
 ttc acc atc agc agt gtg cag gcg gaa gac ctg gca ctt tat tac tgt 336  
 Phe Thr Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Leu Tyr Tyr Cys  
 75 80 85  
 cag caa cat tat agt act cca ttc acg ttc ggc tcg ggg aca aag ttg 384  
 Gln Gln His Tyr Ser Thr Pro Phe Thr Phe Gly Ser Gly Thr Lys Leu  
 90 95 100  
 gaa ata aaa c 394  
 Glu Ile Lys  
 105

<210> 2  
 <211> 131  
 <212> PRT  
 <213> Mouse

<223> Amino acid sequence of L chain V region of mouse  
 anti-HM1.24 antibody

<400> 2  
 Met Gly Phe Lys Met Glu Ser His Phe Leu Val Phe Val Phe Val Phe  
 -20 -15 -10  
 Leu Trp Leu Ser Gly Val Asp Gly Asp Ile Val Met Thr Gln Ser His  
 -5 -1 1 5  
 Lys Phe Met Ser Thr Ser Val Gly Asp Arg Val Ser Ile Thr Cys Lys  
 10 15 20  
 Ala Ser Gln Asp Val Asn Thr Ala Val Ala Trp Tyr Gln Gln Lys Pro  
 25 30 35 40  
 Gly Gln Ser Pro Lys Leu Leu Ile Tyr Ser Ala Ser Asn Arg Tyr Thr  
 45 50 55  
 Gly Val Pro Asp Arg Ile Thr Gly Ser Gly Ser Gly Thr Asp Phe Thr  
 60 65 70  
 Phe Thr Ile Ser Ser Val Gln Ala Glu Asp Leu Ala Leu Tyr Tyr Cys  
 75 80 85  
 Gln Gln His Tyr Ser Thr Pro Phe Thr Phe Gly Ser Gly Thr Lys Leu  
 90 95 100

Glu Ile Lys

105

<210> 3

<211> 418

<212> DNA

<213> Mouse

<223> cDNA coding for H chain V region of mouse anti-HM1.24 antibody

<400> 3

atg gaa tgt aac tgg ata ctt cct ttt att ctg tca gta act tca ggt 48  
Met Glu Cys Asn Trp Ile Leu Pro Phe Ile Leu Ser Val Thr Ser Gly  
-15 -10 -5  
gcc tac tca cag gtt caa ctc cag cag tct ggg gct gag ctg gca aga 96  
Ala Tyr Ser Gln Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg  
-1 1 5 10  
cct ggg gct tca gtg aag ttg tcc tgc aag gct tct ggc tac acc ttt 144  
Pro Gly Ala Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
15 20 25  
act ccc tac tgg atg cag tgg gta aaa cag agg cct gga cag ggt ctg 192  
Thr Pro Tyr Trp Met Gln Trp Val Lys Gln Arg Pro Gly Gln Gly Leu  
30 35 40 45  
gaa tgg att ggg tct att ttt cct gga gat ggt gat act agg tac agt 240  
Glu Trp Ile Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
50 55 60  
cag aag ttc aag ggc aag gcc aca ttg act gca gat aaa tcc tcc agt 288  
Gln Lys Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser  
65 70 75  
aca gcc tac atg caa ctc agc atc ttg gca ttt gag gac tot gcg gtc 336  
Thr Ala Tyr Met Gln Leu Ser Ile Leu Ala Phe Glu Asp Ser Ala Val  
80 85 90  
tat tac tgt gca aga gga tta cga cga ggg ggg tac tac ttt gac tac 384  
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
95 100 105

tgg ggc caa ggc acc act ctc aca gtc tcc tca g  
 Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser  
 110 115 120

<210> 4  
 <211> 139  
 <212> PRT  
 <213> Mouse

<223> Amino acid sequence of H chain V region of mouse  
 anti-HM1.24 antibody

<400> 4  
 Met Glu Cys Asn Trp Ile Leu Pro Phe Ile Leu Ser Val Thr Ser Gly  
 -15 -10 -5  
 Ala Tyr Ser Gln Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg  
 -1 1 5 10  
 Pro Gly Ala Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
 15 20 25  
 Thr Pro Tyr Trp Met Gln Trp Val Lys Gln Arg Pro Gly Gln Gly Leu  
 30 35 40 45  
 Glu Trp Ile Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
 50 55 60  
 Gln Lys Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser  
 65 70 75  
 Thr Ala Tyr Met Gln Leu Ser Ile Leu Ala Phe Glu Asp Ser Ala Val  
 80 85 90  
 Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
 95 100 105  
 Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser  
 110 115 120

<210> 5  
 <211> 11  
 <212> PRT  
 <213> Artificial Sequence

00222E0" 86060560

<220> CDR(1) of L chain V region of anti-HM1.24 antibody  
<223>

<400> 5

Lys Ala Ser Gln Asp Val Asn Thr Ala Val Ala  
1 5 10

<210> 6  
<211> 7  
<212> PRT  
<213> Artificial Sequence

<220> CDR(2) of L chain V region of anti-HM1.24 antibody  
<223>

<400> 6

Ser Ala Ser Asn Arg Tyr Thr  
1 5

<210> 7  
<211> 9  
<212> PRT  
<213> Artificial Sequence

<220> CDR(3) of L chain V region of anti-HM1.24 antibody  
<223>

<400> 7

Gln Gln His Tyr Ser Thr Pro Phe Thr  
1 5

<210> 8  
<211> 5  
<212> PRT  
<213> Artificial Sequence

002220"25050550

<220> CDR(1) of H chain V region of anti-HM1.24 antibody  
<223>

<400> 8  
Pro Tyr Trp Met Gln  
1 5

<210> 9  
<211> 16  
<212> PRT  
<213> Artificial Sequence

<220> CDR(2) of H chain V region of anti-HM1.24 antibody  
<223>

<400> 9  
Ser Ile Phe Gly Asp Gly Asp Thr Arg Tyr Ser Gln Lys Phe Lys Gly  
1 5 10 15

<210> 10  
<211> 11  
<212> PRT  
<213> Artificial Sequence

<220> CDR(3) of H chain V region of anti-HM1.24 antibody  
<223>

<400> 10  
Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
1 5 10

<210> 11  
<211> 379  
<212> DNA  
<213> Artificial Sequence

<220> DNA coding for humanized L chain V region of  
anti-HM1.24 antibody

<223>

<400> 11

atg gga tgg agc tgt atc atc ctc tcc ttg gta gca aca gct aca ggt 48  
Met Gly Trp Ser Cys Ile Ile Leu Ser Leu Val Ala Thr Ala Thr Gly  
-15 -10 -5  
gtc cac tcc gac atc cag atg acc cag agc cca agc agc ctg agc gcc 96  
Val His Ser Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala  
-1 1 5 10  
agc gtg ggt gac aga gtg acc atc acc tgt aag gct agt cag gat gtg 144  
Ser Val Gly Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Val  
15 20 25  
aat act gct gta gcc tgg tac cag cag aag cca gga aag gct cca aag 192  
Asn Thr Ala Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys  
30 35 40 45  
ctg ctg atc tac tgg gca tcc aac cgg tac act ggt gtg cca agc aga 240  
Leu Leu Ile Tyr Ser Ala Ser Asn Arg Tyr Thr Gly Val Pro Ser Arg  
50 55 60  
ttc agc ggt agc ggt agc ggt acc gac ttc acc ttc acc atc agc agc 288  
Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser  
65 70 75  
ctc cag cca gag gac atc gct acc tac tac tgc cag caa cat tat agt 336  
Leu Gln Pro Glu Asp Ile Ala Thr Tyr Tyr Cys Gln Gln His Tyr Ser  
80 85 90  
act cca ttc acg ttc ggc caa ggg acc aag gtg gaa atc aaa c 379  
Thr Pro Phe Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys  
95 100 105

<210> 12

<211> 126

<212> PRT

<213> Artificial Sequence

<220> Humanized L chain V region of anti-HM1.24 antibody

<223>

002220" 86060560

<400> 12

Met Gly Trp Ser Cys Ile Ile Leu Ser Leu Val Ala Thr Ala Thr Gly  
-15 -10 -5  
Val His Ser Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala  
-1 1 5 10  
Ser Val Gly Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Val  
15 20 25  
Asn Thr Ala Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys  
30 35 40 45  
Leu Leu Ile Tyr Ser Ala Ser Asn Arg Tyr Thr Gly Val Pro Ser Arg  
50 55 60  
Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Phe Thr Ile Ser Ser  
65 70 75  
Leu Gln Pro Glu Asp Ile Ala Thr Tyr Tyr Cys Gln Gln His Tyr Ser  
80 85 90  
Thr Pro Phe Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys  
95 100 - 105

<210> 13

<211> 379

<212> DNA

<213> Artificial Sequence

<220> DNA coding for humanized L chain V region of  
anti-HM1.24 antibody

<223>

<400> 13

atg gga tgg agc tgt atc atc ctc tcc ttg gta gca aca gct aca ggt 48  
Met Gly Trp Ser Cys Ile Ile Leu Ser Leu Val Ala Thr Ala Thr Gly  
-15 -10 -5  
gtc cac tcc gac atc cag atg acc cag agc cca agc agc ctg agc gcc 96  
Val His Ser Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala  
-1 1 5 10  
agc gtg ggt gac aga gtg acc atc acc tgt aag gct agt cag gat gtg 144  
Ser Val Gly Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Val  
15 20 25



```

aat act gct gta gcc tgg tac cag cag aag cca gga aag gct cca aag      192
Asn Thr Ala Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys
  30                      35                      40                      45
ctg ctg atc tac tcg gca tcc aac cgg tac act ggt gtg cca agc aga      240
Leu Leu Ile Tyr Ser Ala Ser Asn Arg Tyr Thr Gly Val Pro Ser Arg
                      50                      55                      60
ttc agc ggt agc ggt agt ggt acc gac tac acc ttc acc atc agc agc      288
Phe Ser Gly Ser Gly Ser Gly Thr Asp Tyr Thr Phe Thr Ile Ser Ser
                      65                      70                      75
ctc cag cca gag gac atc gct acc tac tac tgc cag caa cat tat agt      336
Leu Gln Pro Glu Asp Ile Ala Thr Tyr Tyr Cys Gln Gln His Tyr Ser
                      80                      85                      90
act cca ttc acg ttc ggc caa ggg acc aag gtg gaa atc aaa c      379
Thr Pro Phe Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys
                      95                      100                      105

```

<210> 14

<211> 126

<212> PRT

<213> Artificial Sequence

<220> Humanized L chain V region of anti-HM1.24 antibody

<223>

<400> 14

```

Met Gly Trp Ser Cys Ile Ile Leu Ser Leu Val Ala Thr Ala Thr Gly
                      -15                      -10                      -5
Val His Ser Asp Ile Gln Met Thr Gln Ser Pro Ser Ser Leu Ser Ala
                      -1   1                      5                      10
Ser Val Gly Asp Arg Val Thr Ile Thr Cys Lys Ala Ser Gln Asp Val
                      15                      20                      25
Asn Thr Ala Val Ala Trp Tyr Gln Gln Lys Pro Gly Lys Ala Pro Lys
                      30                      35                      40                      45
Leu Leu Ile Tyr Ser Ala Ser Asn Arg Tyr Thr Gly Val Pro Ser Arg
                      50                      55                      60
Phe Ser Gly Ser Gly Ser Gly Thr Asp Tyr Thr Phe Thr Ile Ser Ser
                      65                      70                      75

```

Leu Gln Pro Glu Asp Ile Ala Thr Tyr Tyr Cys Gln Gln His Tyr Ser  
80 85 90  
Thr Pro Phe Thr Phe Gly Gln Gly Thr Lys Val Glu Ile Lys  
95 100 105

<210> 15  
<211> 418  
<212> DNA  
<213> Artificial Sequence

<220> DNA coding for humanized H chain V region(version a)  
of anti-HM1.24 antibody  
<223>

<400> 15  
atg gac tgg acc tgg agg gtc ttc ttc ttg ctg gct gta gct cca ggt 48  
Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
-15 -10 -5  
gct cac tcc cag gtg cag ctg gtg cag tct ggg gct gag gtg aag aag 96  
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
-1 1 5 10  
cct ggg gcc tca gtg aag gtt tcc tgc aag gca tct gga tac acc ttc 144  
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
15 20 25  
act ccc tac tgg atg cag tgg gtg cga cag gcc cct gga caa ggg ctt 192  
Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
30 35 40 45  
gag tgg atg gga tct att ttt cct gga gat ggt gat act agg tac agt 240  
Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
50 55 60  
cag aag ttc aag ggc aga gtc acc atg acc gca gac acg tcc acg agc 288  
Gln Lys Phe Lys Gly Arg Val Thr Met Thr Ala Asp Thr Ser Thr Ser  
65 70 75  
aca gtc tac atg gag ctg agc agc ctg aga tct gag gac acg gcc gtg 336  
Thr Val Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val  
80 85 90

tat tac tgt gcg aga gga tta cga cga ggg ggg tac tac ttt gac tac 384  
 Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
           95                    100                    105

tgg ggg caa ggg acc acg gtc acc gtc tcc tca g 418  
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 110                    115                    120

<210> 16  
 <211> 139  
 <212> PRT  
 <213> Artificial Sequence

<220> Humanized H chain V region(version a) of anti-HM1.24  
 antibody

<223>

<400> 16

Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
                                     -15                                    -10                                    -5  
 Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
                     -1    1                                    5                                    10  
 Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
           15                                    20                                    25  
 Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
           30                                    35                                    40                                    45  
 Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
                                     50                                    55                                    60  
 Gln Lys Phe Lys Gly Arg Val Thr Met Thr Ala Asp Thr Ser Thr Ser  
                                     65                                    70                                    75  
 Thr Val Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val  
           80                                    85                                    90  
 Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
           95                                    100                                    105  
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 110                    115                    120

<210> 17  
 <211> 418

0022220" 86060560

<212> DNA  
 <213> Artificial Sequence  
 <220> DNA coding for humanized H chain V region(version b)  
 of anti-HM1.24 antibody  
 <223>

<400> 17  
 atg gac tgg acc tgg agg gtc ttc ttc ttg ctg gct gta gct cca ggt 48  
 Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
 -15 -10 -5  
 gct cac tcc cag gtg cag ctg gtg cag tct ggg gct gag gtg aag aag 96  
 Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
 -1 1 5 10  
 cct ggg gcc tca gtg aag gtt tcc tgc aag gca tct gga tac acc ttc 144  
 Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
 15 20 25  
 act ccc tac tgg atg cag tgg gtg cga cag gcc cct gga caa ggg ctt 192  
 Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
 30 35 40 45  
 gag tgg atg gga tct att ttt cct gga gat ggt gat act agg tac agt 240  
 Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
 50 55 60  
 cag aag ttc aag ggc aaa gtc acc atg acc gca gac acg tcc acg agc 288  
 Gln Lys Phe Lys Gly Lys Val Thr Met Thr Ala Asp Thr Ser Thr Ser  
 65 70 75  
 aca gtc tac atg gag ctg agc agc ctg aga tct gag gac acg gcc gtg 336  
 Thr Val Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val  
 80 85 90  
 tat tac tgt gcg aga gga tta cga cga ggg ggg tac tac ttt gac tac 384  
 Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
 95 100 105  
 tgg ggg caa ggg acc acg gtc acc gtc tcc tca g 418  
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 110 115 120

<210> 18  
 <211> 139

<212> PRT  
 <213> Artificial Sequence  
 <220> Humanized H chain V region(version b) of anti-HM1.24 antibody  
 <223>

<400> 18  
 Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
 -15 -10 -5  
 Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
 -1 1 5 10  
 Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
 15 20 25  
 Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
 30 35 40 45  
 Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
 50 55 60  
 Gln Lys Phe Lys Gly Lys Val Thr Met Thr Ala Asp Thr Ser Thr Ser  
 65 70 75  
 Thr Val Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val  
 80 85 90  
 Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
 95 100 105  
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 110 115 120

<210> 19  
 <211> 418  
 <212> DNA  
 <213> Artificial Sequence  
 <220> DNA coding for H chain V region(version c) of anti-HM1.24 antibody  
 <223>  
 <400> 19

002220 85060560

```

atg gac tgg acc tgg agg gtc ttc ttc ttg ctg gct gta gct cca ggt      48
Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly
          -15                      -10                      -5

gct cac tcc cag gtg cag ctg gtg cag tct ggg gct gag gtg aag aag      96
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys
          -1   1                      5                      10

cct ggg gcc tca gtg aag gtt tcc tgc aag gca tct gga tac acc ttc      144
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe
          15                      20                      25

act ccc tac tgg atg cag tgg gtg cga cag gcc cct gga caa ggg ctt      192
Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu
          30                      35                      40                      45

gag tgg atg gga tct att ttt cct gga gat ggt gat act agg tac agt      240
Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser
          50                      55                      60

cag aag ttc aag ggc aga gtc act atg acc gca gac aag tcc acg agc      288
Gln Lys Phe Lys Gly Arg Val Thr Met Thr Ala Asp Lys Ser Thr Ser
          65                      70                      75

aca gtc tac atg gag ctg agc agc ctg aga tct gag gac acg gcc gtg      336
Thr Val Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val
          80                      85                      90

tat tac tgt gcg aga gga tta cga cga ggg ggg tac tac ttt gac tac      384
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr
          95                      100                      105

tgg ggg caa ggg acc acg gtc acc gtc tcc tca g                        418
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
          110                      115                      120

```

```

<210>    20
<211>    139
<212>    PRT
<213>    Artificial Sequence

<220>    H chain V region(version c) of anti-HM1.24 antibody
<223>

<400>    20

```

```

Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly
                    -15                -10                -5
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys
        -1   1                5                10
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe
        15                20                25
Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu
        30                35                40                45
Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser
                    50                55                60
Gln Lys Phe Lys Gly Arg Val Thr Met Thr Ala Asp Lys Ser Thr Ser
                    65                70                75
Thr Val Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val
                    80                85                90
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr
                    95                100                105
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
110                115                120

```

```

<210>    21
<211>    418
<212>    DNA
<213>    Artificial Sequence

```

```

<220>    DNA coding for humanized H chain V region(version d)
          of anti-HM1.24 antibody

```

```

<223>

```

```

<400>    21
atg gac tgg acc tgg agg gtc ttc ttc ttg ctg gct gta gct cca ggt      48
Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly
                    -15                -10                -5
gct cac tcc cag gtg cag ctg gtg cag tct ggg gct gag gtg aag aag      96
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys
        -1   1                5                10
cct ggg gcc tca gtg aag gtt tcc tgc aag gca tct gga tac acc ttc      144
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe
        15                20                25

```

act ccc tac tgg atg cag tgg gtg cga cag gcc cct gga caa ggg ctt 192  
 Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
 30 35 40 45  
 gag tgg atg gga tct att ttt cct gga gat ggt gat act agg tac agt 240  
 Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
 50 55 60  
 cag aag ttc aag ggc aaa gtc acc atg acc gca gac aag tcc acg agc 288  
 Gln Lys Phe Lys Gly Lys Val Thr Met Thr Ala Asp Lys Ser Thr Ser  
 65 70 75  
 aca gtc tac atg gag ctg agc agc ctg aga tct gag gac acg gcc gtg 336  
 Thr Val Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val  
 80 85 90  
 tat tac tgt gcg aga gga tta cga cga ggg ggg tac tac ttt gac tac 384  
 Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
 95 100 105  
 tgg ggg caa ggg acc acg gtc acc gtc tcc tca g 418  
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 110 115 120

<210> 22  
 <211> 139  
 <212> PRT  
 <213> Artificial Sequence

<220> H chain V region(version d) of anti-HM1.24 antibody  
 <223>

<400> 22  
 Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
 -15 -10 -5  
 Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
 -1 1 5 10  
 Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
 15 20 25  
 Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
 30 35 40 45  
 Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
 50 55 60



Gln Lys Phe Lys Gly Lys Val Thr Met Thr Ala Asp Lys Ser Thr Ser  
65 70 75  
Thr Val Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val  
80 85 90  
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
95 100 105  
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
110 115 120

<210> 23

<211> 418

<212> DNA

<213> Artificial Sequence

<220> DNA coding for humanized H chain V region(version e)  
of anti-HM1.24 antibody

<223>

<400> 23

atg gac tgg acc tgg agg gtc ttc ttc ttg ctg gct gta gct cca ggt	48
Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly	
-15 -10 -5	
gct cac tcc cag gtg cag ctg gtg cag tct ggg gct gag gtg aag aag	96
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys	
-1 1 5 10	
cct ggg gcc tca gtg aag gtt tcc tgc aag gca tct gga tac acc ttc	144
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe	
15 20 25	
act ccc tac tgg atg cag tgg gtg cga cag gcc cct gga caa ggg ctt	192
Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu	
30 35 40 45	
gag tgg atg gga tct att ttt cct gga gat ggt gat act agg tac agt	240
Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser	
50 55 60	
cag aag ttc aag ggc aga gcc acc ctg acc gca gac acg tcc acg agc	288
Gln Lys Phe Lys Gly Arg Ala Thr Leu Thr Ala Asp Thr Ser Thr Ser	
65 70 75	

aca gtc tac atg gag ctg agc agc ctg aga tct gag gac acg gcc gtg 336  
 Thr Val Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val  
           80                          85                          90  
 tat tac tgt gcg aga gga tta cga cga ggg ggg tac tac ttt gac tac 384  
 Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
           95                          100                         105  
 tgg ggg caa ggg acc acg gtc acc gtc tcc tca g 418  
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 110                          115                         120

<210> 24  
 <211> 239  
 <212> PRT  
 <213> Artificial Sequence

<220> H chain V region(version e) of anti-HM1.24 antibody  
 <223>

<400> 24  
 Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
                           -15                          -10                          -5  
 Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
           -1   1                          5                          10  
 Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
           15                          20                          25  
 Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
           30                          35                          40                          45  
 Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
                           50                          55                          60  
 Gln Lys Phe Lys Gly Arg Ala Thr Leu Thr Ala Asp Thr Ser Thr Ser  
                           65                          70                          75  
 Thr Val Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val  
           80                          85                          90  
 Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
           95                          100                         105  
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 110                          115                         120

Variable	Mean	SD	Min	Max
Age	34.5	10.2	21	55
Gender	0.5	0.5	0	1
Marital status	0.6	0.5	0	1
Education	12.5	1.5	9	16
Income	1500	500	500	3000
Health status	0.7	0.4	0	1
Smoking status	0.3	0.5	0	1
Alcohol consumption	0.2	0.4	0	1
Exercise frequency	0.4	0.5	0	1
Stress level	0.6	0.5	0	1
Sleep quality	0.5	0.5	0	1
Work satisfaction	0.4	0.5	0	1
Life satisfaction	0.5	0.5	0	1
Depression score	10.5	5.0	0	30
Anxiety score	12.0	6.0	0	30
Overall well-being	0.6	0.5	0	1

Variable	Mean	SD	Min	Max
Age	34.5	10.2	21	55
Gender	0.5	0.5	0	1
Marital status	0.6	0.5	0	1
Education	12.5	1.5	9	16
Income	1500	500	500	3000
Health status	0.7	0.4	0	1
Smoking status	0.3	0.5	0	1
Alcohol consumption	0.2	0.4	0	1
Exercise frequency	0.4	0.5	0	1
Stress level	0.6	0.5	0	1
Sleep quality	0.5	0.5	0	1
Work satisfaction	0.4	0.5	0	1
Life satisfaction	0.5	0.5	0	1
Depression score	10.5	5.0	0	30
Anxiety score	12.0	6.0	0	30
Overall well-being	0.6	0.5	0	1

<210> 26  
 <211> 139  
 <212> PRT  
 <213> Artificial Sequence  
  
 <220> Humanized H chain V region(version f) of anti-HM1.24  
 antibody  
 <223>

<400> 26  
 Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
                     -15                    -10                    -5  
 Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
           -1   1                    5                    10  
 Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
       15                    20                    25  
 Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
   30                    35                    40                    45  
 Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
                     50                    55                    60  
 Gln Lys Phe Lys Gly Arg Ala Thr Leu Thr Ala Asp Thr Ser Ser Ser  
                     65                    70                    75  
 Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val  
           80                    85                    90  
 Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
       95                    100                    105  
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 110                    115                    120

<210> 27  
 <211> 418  
 <212> DNA  
 <213> Artificial Sequence  
  
 <220> DNA coding for humanized H chain V region(version g)  
 of anti-HM1.24 antibody  
 <223>

<400> 27

atg gac tgg acc tgg agg gtc ttc ttc ttg ctg gct gta gct cca ggt 48  
Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
-15 -10 -5

gct cac tcc cag gtg cag ctg gtg cag tct ggg gct gag gtg aag aag 96  
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
-1 1 5 10

cct ggg gcc tca gtg aag gtt tcc tgc aag gca tct gga tac acc ttc 144  
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
15 20 25

act ccc tac tgg atg cag tgg gtg cga cag cgc cct gga caa ggg ctt 192  
Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Arg Pro Gly Gln Gly Leu  
30 35 40 45

gag tgg atg gga tct att ttt cct gga gat ggt gat act agg tac agt 240  
Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
50 - 55 60

cag aag ttc aag ggc aga gtc acc atg acc gca gac acg tcc acg agc 288  
Gln Lys Phe Lys Gly Arg Val Thr Met Thr Ala Asp Thr Ser Thr Ser  
65 70 75

aca gtc tac atg gag ctg agc agc ctg aga tct gag gac acg gcc gtg 336  
Thr Val Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val  
80 85 90

tat tac tgt gcg aga gga tta cga cga ggg ggg tac tac ttt gac tac 384  
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
95 100 105

tgg ggg caa ggg acc acg gtc acc gtc tcc tca g 418  
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
110 115 120

<210> 28

<211> 139

<212> PRT

<213> Artificial Sequence

<220> DNA coding for humanized H chain V region(version g)  
of anti-HM1.24 antibody

<223>

&lt;400&gt; 28

```

Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly
                    -15                -10                -5
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys
        -1  1                      5                      10
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe
        15                      20                      25
Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Arg Pro Gly Gln Gly Leu
        30                      35                      40                      45
Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser
                    50                      55                      60
Gln Lys Phe Lys Gly Arg Val Thr Met Thr Ala Asp Thr Ser Thr Ser
                    65                      70                      75
Thr Val Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val
                    80                      85                      90
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr
                    95                      100                      105
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
110                      115                      120

```

&lt;210&gt; 29

&lt;211&gt; 418

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

<220> DNA coding for humanized H chain V region(version h)  
of anti-HM1.24 antibody

&lt;223&gt;

&lt;400&gt; 29

```

atg gac tgg acc tgg agg gtc ttc ttc ttg ctg gct gta gct cca ggt      48
Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly
                    -15                -10                -5
gct cac tcc cag gtg cag ctg gtg cag tct ggg gct gag gtg aag aag      96
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys
        -1  1                      5                      10

```

```

cct ggg gcc tca gtg aag gtt tcc tgc aag gca tct gga tac acc ttc      144
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe
      15              20              25

act ccc tac tgg atg cag tgg gtg cga cag gcc cct gga caa ggg ctt      192
Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu
      30              35              40              45

gag tgg atg gga tct att ttt cct gga gat ggt gat act agg tac agt      240
Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser
              50              55              60

cag aag ttc aag ggc aaa gtc acc atg acc gca gac acg tcc tcg agc      288
Gln Lys Phe Lys Gly Lys Val Thr Met Thr Ala Asp Thr Ser Ser Ser
              65              70              75

aca gcc tac atg gag ctg agc agc ctg aga tct gag gac acg gcc gtg      336
Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val
              80              85              90

tat tac tgt gcg aga gga tta cga cga ggg ggg tac tac ttt gac tac      384
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr
              95              100              105

tgg ggg caa ggg acc acg gtc acc gtc tcc tca g      418
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
110              115              120

```

```

<210> 30
<211> 239
<212> PRT
<213> Artificer Sequence

```

```

<220> Humanized H chain V region(version h) of anti-HM1.24
      antibody

```

```

<223>

```

```

<400> 30
Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly
              -15              -10              -5
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys
              -1  1              5              10
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe
      15              20              25

```

Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
 30 35 40 45  
 Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
 50 55 60  
 Gln Lys Phe Lys Gly Lys Val Thr Met Thr Ala Asp Thr Ser Ser Ser  
 65 70 75  
 Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val  
 80 85 90  
 Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
 95 100 105  
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 110 115 120

<210> 31  
 <211> 418  
 <212> DNA  
 <213> Artificial Sequence -

<220> DNA coding for humanized H chain V region(version i)  
 of anti-HM1.24 antibody

<223>

<400> 31

atg gac tgg acc tgg agg gtc ttc ttc ttg ctg gct gta gct oca ggt 48  
 Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
 -15 -10 -5  
 gct cac tcc cag gtg cag ctg gtg cag tct ggg gct gag gtg aag aag 96  
 Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
 -1 1 5 10  
 cct ggg gcc tca gtg aag gtt tcc tgc aag gca tct gga tac acc ttc 144  
 Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
 15 20 25  
 act ccc tac tgg atg cag tgg gtg cga cag gcc cct gga caa ggg ctt 192  
 Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
 30 35 40 45  
 gag tgg atg gga tct att ttt cct gga gat ggt gat act agg tac agt 240  
 Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
 50 55 60



cag aag ttc aag ggc aaa gtc acc atg acc gca gac acg tcc tcg agc 288  
 Gln Lys Phe Lys Gly Lys Val Thr Met Thr Ala Asp Thr Ser Ser Ser  
           65                          70                          75  
 aca gcc tac atg gag ctg agc agc ctg gca ttt gag gac acg gcc gtg 336  
 Thr Ala Tyr Met Glu Leu Ser Ser Leu Ala Phe Glu Asp Thr Ala Val  
           80                          85                          90  
 tat tac tgt gcg aga gga tta cga cga ggg ggg tac tac ttt gac tac 384  
 Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
           95                          100                         105  
 tgg ggg caa ggg acc acg gtc acc gtc tcc tca g 418  
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 110                          115                          120

<210> 32

<211> 139

<212> PRT

<213> Artificial Sequence -

<220> Humanized H chain V region(version i) of anti-HM1.24 antibody

<223>

<400> 32

Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
                           -15                          -10                          -5  
 Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
           -1   1                          5                          10  
 Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
           15                          20                          25  
 Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
           30                          35                          40                          45  
 Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
                           50                          55                          60  
 Gln Lys Phe Lys Gly Lys Val Thr Met Thr Ala Asp Thr Ser Ser Ser  
           65                          70                          75  
 Thr Ala Tyr Met Glu Leu Ser Ser Leu Ala Phe Glu Asp Thr Ala Val  
           80                          85                          90



tat tac tgt gcg aga gga tta cga cga ggg ggg tac tac ttt gac tac 384  
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
95 100 105  
tgg ggg caa ggg acc acg gtc acc gtc tcc tca g 418  
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
110 115 120

<210> 34  
<211> 139  
<212> PRT  
<213> Artificial Sequence

<220> Humanized H chain V region(version j) of anti-HM1.24  
antibody

<223>

<400> 34

Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
-15 -10 -5  
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
-1 1 5 10  
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
15 20 25  
Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
30 35 40 45  
Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
50 55 60  
Gln Lys Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Thr Ser Ser Ser  
65 70 75  
Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val  
80 85 90  
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
95 100 105  
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
110 115 120

<210> 35  
<211> 418

<212> DNA  
 <213> Artificial Sequence  
 <220> DNA coding for H chain V region(version k) of  
 anti-HM1.24 antibody  
 <223>

<400> 35  
 atg gac tgg acc tgg agg gtc ttc ttc ttg ctg gct gta gct cca ggt 48  
 Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
 -15 -10 -5  
 gct cac tcc cag gtg cag ctg gtg cag tct ggg gct gag gtg aag aag 96  
 Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
 -1 1 5 10  
 cct ggg gcc tca gtg aag gtt tcc tgc aag gca tct gga tac acc ttc 144  
 Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
 15 20 25  
 act ccc tac tgg atg cag tgg gtg cga cag gcc cct gga caa ggg ctt 192  
 Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
 30 35 40 45  
 gag tgg atg gga tct att ttt cct gga gat ggt gat act agg tac agt 240  
 Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
 50 55 60  
 cag aag ttc aag ggc aaa gtc acc atg acc gca gac acg tcc tcg agc 288  
 Gln Lys Phe Lys Gly Lys Val Thr Met Thr Ala Asp Thr Ser Ser Ser  
 65 70 75  
 aca gcc tac atg cag ctg agc agc cta aga tct gag gac acg gcc gtg 336  
 Thr Ala Tyr Met Gln Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val  
 80 85 90  
 tat tac tgt gcg aga gga tta cga cga ggg ggg tac tac ttt gac tac 384  
 Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
 95 100 105  
 tgg ggg caa ggg acc acg gtc acc gtc tcc tca g 418  
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 110 115 120

<210> 36  
 <211> 139

<212> PRT  
 <213> Artificial Seam  
 <220> Humanized H chain V region(version k) of anti-HM1.24 antibody  
 <223>

<400> 36  
 Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
                   -15                  -10                  -5  
 Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
           -1   1                  5                  10  
 Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
       15                  20                  25  
 Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
   30                  35                  40                  45  
 Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
                   50                  55                  60  
 Gln Lys Phe Lys Gly Lys Val Thr Met Thr Ala Asp Thr Ser Ser Ser  
           65                  70                  75  
 Thr Ala Tyr Met Gln Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val  
       80                  85                  90  
 Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
       95                  100                  105  
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 110                  115                  120

<210> 37  
 <211> 418  
 <212> DNA  
 <213> Artificial Sequence  
 <220> DNA coding for humanized H chain V region(version l)  
           of anti-HM1.24 antibody  
 <223>  
 <400> 37

002220" 26060560

002220"86060560

```

atg gac tgg acc tgg agg gtc ttc ttc ttg ctg gct gta gct cca ggt      48
Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly
          -15                -10                -5

gct cac tcc cag gtg cag ctg gtg cag tct ggg gct gag gtg aag aag      96
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys
          -1   1                5                10

cct ggg gcc tca gtg aag gtt tcc tgc aag gca tct gga tac acc ttc      144
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe
          15                20                25

act ccc tac tgg atg cag tgg gtg cga cag gcc cct gga caa ggg ctt      192
Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu
          30                35                40                45

gag tgg atg gga tct att ttt cct gga gat ggt gat act agg tac agt      240
Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser
          50                55                60

cag aag ttc aag ggc aaa gtc acc atg acc gca gac acg tcc tcg agc      288
Gln Lys Phe Lys Gly Lys Val Thr Met Thr Ala Asp Thr Ser Ser Ser
          65                70                75

aca gcc tac atg cag ctg agc atc ctg aga tct gag gac acg gcc gtg      336
Thr Ala Tyr Met Gln Leu Ser Ile Leu Arg Ser Glu Asp Thr Ala Val
          80                85                90

tat tac tgt gcg aga gga tta cga cga ggg ggg tac tac ttt gac tac      384
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr
          95                100                105

tgg ggg caa ggg acc acg gtc acc gtc tcc tca g      418
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
          110                115                120

```

<210> 38  
 <211> 139  
 <212> PRT  
 <213> Artificial Sequence

<220> Humanized H chain V region(version 1) of anti-HM1.24  
 antibody

<223>

<400> 38

Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
 -15 -10 -5  
 Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
 -1 1 5 10  
 Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
 15 20 25  
 Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
 30 35 40 45  
 Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
 50 55 60  
 Gln Lys Phe Lys Gly Lys Val Thr Met Thr Ala Asp Thr Ser Ser Ser  
 65 70 75  
 Thr Ala Tyr Met Gln Leu Ser Ile Leu Arg Ser Glu Asp Thr Ala Val  
 80 85 90  
 Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
 95 100 105  
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 110 115 120

<210> 39  
 <211> 418  
 <212> DNA  
 <213> Artificial Sequence  
 <220> DNA coding for humanized H chain V region(version m)  
 of anti-HM1.24 antibody  
 <223>

<400> 39  
 atg gac tgg acc tgg agg gtc ttc ttc ttg ctg gct gta gct cca ggt 48  
 Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
 -15 -10 -5  
 gct cac tcc cag gtg cag ctg gtg cag tct ggg gct gag gtg aag aag 96  
 Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
 -1 1 5 10  
 cct ggg gcc tca gtg aag gtt tcc tgc aag gca tct gga tac acc ttc 144  
 Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
 15 20 25

act ccc tac tgg atg cag tgg gtg cga cag gcc cct gga caa ggg ctt 192  
 Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
 30 35 40 45  
 gag tgg atg gga tct att ttt cct gga gat ggt gat act agg tac agt 240  
 Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
 50 55 60  
 cag aag ttc aag ggc aaa gtc acc atg acc gca gac acg tcc tcg agc 288  
 Gln Lys Phe Lys Gly Lys Val Thr Met Thr Ala Asp Thr Ser Ser Ser  
 65 70 75  
 aca gcc tac atg cag ctg agc atc ctg aga tct gag gac tcg gcc gtg 336  
 Thr Ala Tyr Met Gln Leu Ser Ile Leu Arg Ser Glu Asp Ser Ala Val  
 80 85 90  
 tat tac tgt gcg aga gga tta cga cga ggg ggg tac tac ttt gac tac 384  
 Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
 95 100 105  
 tgg ggg caa ggg acc acg gtc acc gtc tcc tca g 418  
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 110 115 120

<210> 40  
 <211> 139  
 <212> PRT  
 <213> Artificial Sequence

<220> Humanized H chain V region(version m) of anti-HM1.24  
 antibody

<223>

<400> 40

Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
 -15 -10 -5  
 Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
 -1 1 5 10  
 Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
 15 20 25  
 Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
 30 35 40 45



Glu	Trp	Met	Gly	Ser	Ile	Phe	Pro	Gly	Asp	Gly	Asp	Thr	Arg	Tyr	Ser
				50				55				60			
Gln	Lys	Phe	Lys	Gly	Lys	Val	Thr	Met	Thr	Ala	Asp	Thr	Ser	Ser	Ser
				65				70				75			
Thr	Ala	Tyr	Met	Gln	Leu	Ser	Ile	Leu	Arg	Ser	Glu	Asp	Ser	Ala	Val
				80				85				90			
Tyr	Tyr	Cys	Ala	Arg	Gly	Leu	Arg	Arg	Gly	Gly	Tyr	Tyr	Phe	Asp	Tyr
				95				100				105			
Trp	Gly	Gln	Gly	Thr	Thr	Val	Thr	Val	Ser	Ser					
				110				115				120			

<210> 41

<212> DNA

<220> DNA coding for humanized H chain V region(version n)  
of anti-HM1.24 antibody

<400> 41

Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly

gct cac tcc caq gtg caq ctg gtg caq tct ggg gct gag gtg aag aag 96

-1      1                          5                          10

Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe

act ccc tac tgg atg cag tgg gtg cga cag gcc cct gga caa ggg ctt 192

30                      35                      40                      45

Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser



Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
 95 100 105  
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 110 115 120

<210> 43  
 <211> 418  
 <212> DNA  
 <213> Artificial Sequence

<220> DNA coding for humanized H chain V region(version o)  
 of anti-HM1.24 antibody

<223>

<400> 43  
 atg gac tgg acc tgg agg gtc ttc ttc ttg ctg gct gta gct cca ggt 48  
 Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
 -15 -10 -5  
 gct cac tcc cag gtg cag ctg gtg cag tct ggg gct gag gtg aag aag 96  
 Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
 -1 1 5 10  
 cct ggg gcc tca gtg aag gtt tcc tgc aag gca tct gga tac acc ttc 144  
 Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
 15 20 25  
 act ccc tac tgg atg cag tgg gtg cga cag gcc cct gga caa ggg ctt 192  
 Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
 30 35 40 45  
 gag tgg atg gga tct att ttt cct gga gat ggt gat act agg tac agt 240  
 Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
 50 55 60  
 cag aag ttc aag ggc aaa gtc acc atg acc gca gac acg tcc tcg agc 288  
 Gln Lys Phe Lys Gly Lys Val Thr Met Thr Ala Asp Thr Ser Ser Ser  
 65 70 75  
 aca gcc tac atg gag ctg agc agc ctg aga tct gag gac tcg gcc gta 336  
 Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Ser Ala Val  
 80 85 90

002220" 86060560





<212> PRT  
 <213> Artificial Sequence  
 <220> Humanized H chain V region(version p) of anti-HM1.24 antibody  
 <223>

<400> 46  
 Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
 -15 -10 -5  
 Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
 -1 1 5 10  
 Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
 15 20 25  
 Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
 30 35 40 45  
 Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
 50 55 60  
 Gln Lys Phe Lys Gly Arg Val Thr Met Thr Ala Asp Thr Ser Thr Ser  
 65 70 75  
 Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val  
 80 85 90  
 Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
 95 100 105  
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 110 115 120

<210> 47  
 <211> 418  
 <212> DNA  
 <213> Artificial Sequence  
 <220> DNA coding for humanized H chain V region(version p) of anti-HM1.24 antibody  
 <223>  
 <400> 47

atg gac tgg acc tgg agg gtc ttc ttc ttg ctg gct gta gct cca ggt	48
Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly	
-15 -10 -5	
gct cac tcc cag gtg cag ctg gtg cag tct ggg gct gag gtg aag aag	96
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys	
-1 1 5 10	
cct ggg gcc tca gtg aag gtt tcc tgc aag gca tct gga tac acc ttc	144
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe	
15 20 25	
act ccc tac tgg atg cag tgg gtg cga cag gcc cct gga caa ggg ctt	192
Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu	
30 35 40 45	
gag tgg atg gga tct att ttt cct gga gat ggt gat act agg tac agt	240
Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser	
50 55 60	
cag aag ttc aag ggc aga gtc acc atg acc gca gac acg tcc tcg agc	288
Gln Lys Phe Lys Gly Arg Val Thr Met Thr Ala Asp Thr Ser Ser Ser	
65 70 75	
aca gtc tac atg gag ctg agc agc ctg aga tct gag gac acg gcc gtg	336
Thr Val Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val	
80 85 90	
tat tac tgt gcg aga gga tta cga cga ggg ggg tac tac ttt gac tac	384
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr	
95 100 105	
tgg ggg caa ggg acc acg gtc acc gtc tcc tca g	418
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser	
110 115 120	

<210> 48

<211> 139

<212> PRT

<213> Artificial Sequence

<220> Humanized H chain V region(version p) of anti-HM1.24 antibody

<223>

<400> 48

Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
-15 -10 -5  
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
-1 1 5 10  
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
15 20 25  
Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
30 35 40 45  
Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
50 55 60  
Gln Lys Phe Lys Gly Arg Val Thr Met Thr Ala Asp Thr Ser Ser Ser  
65 70 75  
Thr Val Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val  
80 85 90  
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
95 100 105  
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
110 115 120

<210> 49  
<211> 418  
<212> DNA  
<213> Artificial Sequence

<220> DNA coding for humanized H chain V region(version r)  
of anti-HM1.24 antibody

<223>

<400> 49  
atg gac tgg acc tgg agg gtc ttc ttc ttg ctg gct gta gct cca ggt 48  
Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
-15 -10 -5  
gct cac tcc cag gtg cag ctg gtg cag tct ggg gct gag gtg aag aag 96  
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
-1 1 5 10  
cct ggg gcc tca gtg aag gtt tcc tgc aag gca tct gga tac acc ttc 144  
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
15 20 25



act ccc tac tgg atg cag tgg gtg cga cag gcc cct gga caa ggg ctt 192  
 Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
 30 35 40 45  
 gag tgg atg gga tct att ttt cct gga gat ggt gat act agg tac agt 240  
 Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
 50 55 60  
 cag aag ttc aag ggc aga gtc acc atg acc gca gac aag tcc acg agc 288  
 Gln Lys Phe Lys Gly Arg Val Thr Met Thr Ala Asp Lys Ser Thr Ser  
 65 70 75  
 aca gcc tac atg gag ctg agc agc ctg aga tct gag gac acg gcc gtg 336  
 Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val  
 80 85 90  
 tat tac tgt gcg aga gga tta cga cga ggg ggg tac tac ttt gac tac 384  
 Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
 95 100 105  
 tgg ggg caa ggg acc acg gtc acc gtc tcc tca g 418  
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 110 115 120

<210> 50  
 <211> 139  
 <212> PRT  
 <213> Artificial Sequence

<220> Humanized H chain V region(version r) of anti-HM1.24  
 antibody

<223>

<400> 50  
 Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
 -15 -10 -5  
 Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
 -1 1 5 10  
 Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
 15 20 25  
 Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
 30 35 40 45

Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
                             50                            55                            60  
 Gln Lys Phe Lys Gly Arg Val Thr Met Thr Ala Asp Lys Ser Thr Ser  
                             65                            70                            75  
 Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val  
                             80                            85                            90  
 Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
                             95                            100                            105  
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 110                            115                            120

<210> 51  
 <211> 40  
 <212> DNA  
 <213> Artificial Sequence

<220> Primer  
 <223> Synthetic DNA

<400> 51  
 actagtcgac atgaagttgc ctgttaggct gttggtgctg

40

<210> 52  
 <211> 39  
 <212> DNA  
 <213> Artificial Sequence

<220> Primer  
 <223> Synthetic DNA

<400> 52  
 actagtcgac atggagwcag acacactcct gytatgggt

39

<210> 53  
 <211> 40  
 <212> DNA  
 <213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 53

actagtcgac atgagtggtgc tcaactcaggt cctggsgttg

40

<210> 54

<211> 43

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 54

actagtcgac atgaggrccc ctgctcagwt tyttggmwtc ttg

43

<210> 55

<211> 40

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 55

actagtcgac atggatttwc aggtgcagat twtcagcttc

40

<210> 56

<211> 37

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA



002220"26060560

<210> 60  
<211> 37  
<212> DNA  
<213> Artificial Sequence

<220> Primer  
<223> Synthetic DNA

<400> 60  
actagtcgac atgtatatat gtttgttgtc tatttct

37

<210> 61  
<211> 38  
<212> DNA  
<213> Artificial Sequence

<220> Primer  
<223> Synthetic DNA

<400> 61  
actagtcgac atggaagccc cagctcagct tctttcc

38

<210> 62  
<211> 27  
<212> DNA  
<213> Artificial Sequence

<220> Primer  
<223> Synthetic DNA

<400> 62  
ggatcccggtg tggatggtgg gaagatg

27

<210> 63  
<211> 25

00222E0" 86060560

<212> DNA  
<213> Artificial Sequence

<220> Primer  
<223> Synthetic DNA

<400> 63  
tagagtcacc gaggagccag ttgta

25

<210> 64  
<211> 26  
<212> DNA  
<213> Artificial Sequence

<220> Primer  
<223> Synthetic DNA

<400> 64  
ggatccoggg agtggataga ccgatg

26

<210> 65  
<211> 34  
<212> DNA  
<213> Artificial Sequence

<220> Primer  
<223> Synthetic DNA

<400> 65  
gataagcttc caccatgggc ttcaagatgg agtc

34

<210> 66  
<211> 34  
<212> DNA  
<213> Artificial Sequence

00222ED"86060560

<220> Primer  
<223> Synthetic DNA

<400> 66  
gataagcttc caccatggaa tgtaactgga tact

34

<210> 67  
<211> 34  
<212> DNA  
<213> Artificial Sequence

<220> Primer  
<223> Synthetic DNA

<400> 67  
ggcggatcca ctcacgtttt atttccaact ttgt

34

<210> 68  
<211> 34  
<212> DNA  
<213> Artificial Sequence

<220> Primer  
<223> Synthetic DNA

<400> 68  
ggcggatcca ctcacctgag gagactgtga gagt

34

<210> 69  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220> Primer  
<223> Synthetic DNA

<400> 69

cagacagtgg ttcaaagt

18

<210> 70

<211> 26

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 70

gaattcggat ccactcacgt ttgatt

26

<210> 71

<211> 48

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 71

agtcaggatg tgaatactgc tgtagcctgg taccagcaga agccagga

48

<210> 72

<211> 39

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 72

gcattccaacc ggtacactgg tgtgccaaagc agattcagc

39

0022220"26060560



<210> 73  
 <211> 45  
 <212> DNA  
 <213> Artificial Sequence

<220> Primer  
 <223> Synthetic DNA

<400> 73  
 caacattata gtactccatt cacgttcggc caagggacca aggtg

45

<210> 74  
 <211> 47  
 <212> DNA  
 <213> Artificial Sequence

<220> Primer  
 <223> Synthetic DNA

<400> 74  
 gcagatttca catcctgact ggccttacag gtgatgggtca ctctgtc

47

<210> 75  
 <211> 38  
 <212> DNA  
 <213> Artificial Sequence

<220> Primer  
 <223> Synthetic DNA

<400> 75  
 acaccagtgt accggttgga tgccgagtag atcagcag

38

<210> 76  
 <211> 41  
 <212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 76

gtgaatggag tactataatg ttgctggcag tagtaggtag c

41

<210> 77

<211> 31

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 77

ggtaccgact acaccttcac catcagcagc c

31

<210> 78

<211> 31

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 78

ggtgaagggtg tagtcggtac cgctaccgct a

31

<210> 79

<211> 144

<212> DNA

<213> Artificial Sequence

<220> Primer

002250" 86060560

<223> Synthetic DNA

<400> 79

atgccttgca ggaaaccttc actgaggccc caggettott cacctcagcc coagactgca 60  
ccagctgcac ctgggagtga gcacctggag ctacagccag caagaagaag acctccagg 120  
tccagtccat ggtggaagct tate 144

<210> 80

<211> 130

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 80

tcagtgaagg tttcctgcaa ggcattctgga tacaccttca ctccctactg gatgcagtgg 60  
gtgacgacagg cccctggaca agggcttgag tggatgggat ctatttttcc tggagatggt 120  
gataactaggt 130

<210> 81

<211> 131

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 81

aatacacggc cgtgtcctca gatctcaggc tgctcagctc catgtagact gtgctcgtgg 60  
acgtgtctgc ggtcatggtg actctgccct tgaacttotg actgtaccta gtatcaccat 120  
ctccaggaaa a 131

<210> 82

<211> 119

<212> DNA

<213> Artificial Sequence

0022E0" 85050550

<220> Primer

<223> Synthetic DNA

<400> 82

gagatctgag gacacggccg tgtattactg tgcgagagga ttacgacgag ggggggtacta 60  
ctttgactac tgggggcaag ggaccacggt caccgtctcc tcagggtgagt ggatccgac 119

<210> 83

<211> 25

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 83

gataagcttc caccatggac tggac 25

<210> 84

<211> 25

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 84

gtcggatcca ctcacctgag gagac 25

<210> 85

<211> 26

<212> DNA

<213> Artificial Sequence

<220> Primer

002220 86060560

<223> Synthetic DNA

<400> 85

aagttcaagg gcaaagtcac catgac

26

<210> 86

<211> 26

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 86

gtcatggtga ctttgcctt gaactt

26

<210> 87

<211> 26

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 87

atgaccgcag acaagtccac gagcac

26

<210> 88

<211> 26

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 88

gtgctcgtgg acttgtctgc ggtcat

26

<210> 89

<211> 47

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 89

aagtccaagg gcaaagtcac catgaccgca gacaagtcca cgagcac

47

<210> 90

<211> 47

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 90

gtgctcgtgg acttgtctgc ggtcatggtg actttgcct tgaactt

47

<210> 91

<211> 38

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 91

aagtccaagg gcagagccac cctgaccgca gacacgtc

38

<210> 92

0022E0" 85060560

<211> 38  
<212> DNA  
<213> Artificial Sequence

<220> Primer  
<223> Synthetic DNA

<400> 92  
gacgtgtctg cggtcagggt ggctctgccc ttgaactt

38

<210> 93  
<211> 18  
<212> DNA  
<213> Artificial Sequence

<220> Primer  
<223> Synthetic DNA

<400> 93  
cagacagtgg ttcaaagt

18

<210> 94  
<211> 17  
<212> DNA  
<213> Artificial Sequence

<220> Primer  
<223> Synthetic DNA

<400> 94  
gccccaaagc caaggtc

17

<210> 95  
<211> 23  
<212> DNA  
<213> Artificial Sequence

0022220" 86060560

<220> Primer  
<223> Synthetic DNA

<400> 95  
atttttcctg gagatggtga tac

23

<210> 96  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220> Primer  
<223> Synthetic DNA

<400> 96  
gtatcaccat ctccaggaaa tat

23

<210> 97  
<211> 418  
<212> DNA  
<213> Artificial Sequence

<220> DNA coding for humanized H chain V region(native/ver  
sion a mix) of anti-HM1.24 antibody  
<223>

<400> 97  
atg gaa tgt aac tgg ata ctt cct ttt att ctg tca gta act tca ggt  
Met Glu Cys Asn Trp Ile Leu Pro Phe Ile Leu Ser Val Thr Ser Gly  
-15 -10 -5  
gcc tac tca cag gtt caa ctc cag cag tct ggg gct gag ctg gca aga  
Ala Tyr Ser Gln Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg  
-1 1 5 10

48

96



0022220" 86060560

```

cct ggg gct tca gtg aag ttg tcc tgc aag gct tct ggc tac acc ttt      144
Pro Gly Ala Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe
      15              20              25

act ccc tac tgg atg cag tgg gta aaa cag agg cct gga cag ggt ctg      192
Thr Pro Tyr Trp Met Gln Trp Val Lys Gln Arg Pro Gly Gln Gly Leu
      30              35              40              45

gaa tgg att ggg tct att ttt cct gga gat ggt gat act agg tac agt      240
Glu Trp Ile Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser
              50              55              60

cag aag ttc aag ggc aga gtc acc atg acc gca gac acg tcc acg agc      288
Gln Lys Phe Lys Gly Arg Val Thr Met Thr Ala Asp Thr Ser Thr Ser
              65              70              75

aca gtc tac atg gag ctg agc agc ctg aga tct gag gac acg gcc gtg      336
Thr Val Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val
              80              85              90

tat tac tgt gcg aga gga tta cga cga ggg ggg tac tac ttt gac tac      384
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr
              95              100              105

tgg ggg caa ggg acc acg gtc acc gtc tcc tca g      418
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
110              115              120

```

<210> 98  
 <211> 139  
 <212> PRT  
 <213> Artificial Sequence

<220> Humanized H chain V region(native/version a mix) of  
 anti-HM1.24 antibody

<223>

<400> 98  
 Met Glu Cys Asn Trp Ile Leu Pro Phe Ile Leu Ser Val Thr Ser Gly  
 -15 -10 -5  
 Ala Tyr Ser Gln Val Gln Leu Gln Gln Ser Gly Ala Glu Leu Ala Arg  
 -1 1 5 10  
 Pro Gly Ala Ser Val Lys Leu Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
 15 20 25

Thr Pro Tyr Trp Met Gln Trp Val Lys Gln Arg Pro Gly Gln Gly Leu  
 30 35 40 45  
 Glu Trp Ile Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
 50 55 60  
 Gln Lys Phe Lys Gly Arg Val Thr Met Thr Ala Asp Thr Ser Thr Ser  
 65 70 75  
 Thr Val Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val  
 80 85 90  
 Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
 95 100 105  
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 110 115 120

<210> 99

<211> 418

<212> DNA

<213> Artificial Sequence -

<220> DNA coding for humanized C chain V region(native/ver  
 sion a mix) of anti-HM1.24 antibody

<223>

<400> 99

atg gac tgg acc tgg agg gtc ttc ttc ttg ctg gct gta gct cca ggt 48  
 Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
 -15 -10 -5  
 gct cac tcc cag gtg cag ctg gtg cag tct ggg gct gag gtg aag aag 96  
 Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
 -1 1 5 10  
 cct ggg gcc tca gtg aag gtt tcc tgc aag gca tct gga tac acc ttc 144  
 Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
 15 20 25  
 act ccc tac tgg atg cag tgg gtg cga cag gcc cct gga caa ggg ctt 192  
 Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
 30 35 40 45  
 gag tgg atg gga tct att ttt cct gga gat ggt gat act agg tac agt 240  
 Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
 50 55 60

002220" 85050560

```

cag aag ttc aag ggc aag gcc aca ttg act gca gat aaa tcc tcc agt      288
Gln Lys Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser
      65              70              75

aca gcc tac atg caa ctc agc atc ttg gca ttt gag gac tct gcg gtc      336
Thr Ala Tyr Met Gln Leu Ser Ile Leu Ala Phe Glu Asp Ser Ala Val
      80              85              90

tat tac tgt gca aga gga tta cga cga ggg ggg tac tac ttt gac tac      384
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr
      95              100             105

tgg ggc caa ggc acc act ctc aca gtc tcc tca g      418
Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser
     110              115              120

```

<210> 100

<211> 139

<212> PRT

<213> Artificial Sequence -

<220> Humanized C chain V region(native/version a mix) of  
anti-HM1.24 antibody

<223>

<400> 100

```

Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly
      -15              -10              -5

Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys
      -1  1              5              10

Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe
      15              20              25

Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu
      30              35              40              45

Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser
      50              55              60

Gln Lys Phe Lys Gly Lys Ala Thr Leu Thr Ala Asp Lys Ser Ser Ser
      65              70              75

Thr Ala Tyr Met Gln Leu Ser Ile Leu Ala Phe Glu Asp Ser Ala Val
      80              85              90

```

Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
 95 100 105  
 Trp Gly Gln Gly Thr Thr Leu Thr Val Ser Ser  
 110 115 120

<210> 101  
 <211> 38  
 <212> DNA  
 <213> Artificial Sequence

<220> Primer  
 <223> Synthetic DNA

<400> 101  
 ctggttcggc ccacctctga aggttcaga atcgatag

38

<210> 102  
 <211> 35  
 <212> DNA  
 <213> Artificial Sequence

<220> Primer  
 <223> Synthetic DNA

<400> 102  
 gcagacacgt cctogagcac agcctacatg gagct

35

<210> 103  
 <211> 35  
 <212> DNA  
 <213> Artificial Sequence

<220> Primer  
 <223> Synthetic DNA

<400> 103  
 agctccatgt aggtgtgtct cgaggacgtg tctgc

35

<210> 104  
 <211> 26  
 <212> DNA  
 <213> Artificial Sequence

<220> Primer  
 <223> Synthetic DNA

<400> 104  
 tgggtgcgac agcgccctgg acaagg

26

<210> 105  
 <211> 26  
 <212> DNA  
 <213> Artificial Sequence

<220> Primer  
 <223> Synthetic DNA

<400> 105  
 ccttgtccag ggcgctgtcg caccca

26

<210> 106  
 <211> 41  
 <212> DNA  
 <213> Artificial Sequence

<220> Primer  
 <223> Synthetic DNA

<400> 106  
 tacatggagc tgagcagcct ggcatttgag gacacggccg t

41

<210> 107  
 <211> 41



<220> Primer  
<223> Synthetic DNA

<400> 110  
gcctacatgc agctgagcag cct

23

<210> 111  
<211> 23  
<212> DNA  
<213> Artificial Sequence

<220> Primer  
<223> Synthetic DNA

<400> 111  
aggctgctca gctgcatgta ggc

23

<210> 112  
<211> 38  
<212> DNA  
<213> Artificial Sequence

<220> Primer  
<223> Synthetic DNA

<400> 112  
gcctacatgc agctgagcat cctgagatct gaggacac

38

<210> 113  
<211> 35  
<212> DNA  
<213> Artificial Sequence

<220> Primer  
<223> Synthetic DNA

<400> 113  
gatctcagga tgctcagctg catgtaggct gtgct

35

<210> 114  
<211> 50  
<212> DNA  
<213> Artificial Sequence

<220> Primer  
<223> Synthetic DNA

<400> 114  
gcctacatgc agctgagcat cctgagatct gaggactcgg ccgtgtatta

50

<210> 115  
<211> 50  
<212> DNA  
<213> Artificial Sequence

<220> Primer  
<223> Synthetic DNA

<400> 115  
acggcgaggt cctcagatct caggatgctc agctgcatgt aggtgtgct

50

<210> 116  
<211> 20  
<212> DNA  
<213> Artificial Sequence

<220> Primer  
<223> Synthetic DNA

<400> 116  
gagctgagca tootgagatc

20





<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 120

gcagacacgt ccacgagcac agcctacatg gagct

35

<210> 121

<211> 35

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 121

agctccatgt aggctgtgct cgtggacgtg tctgc

35

<210> 122

<211> 35

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 122

gcagacaagt cctcgagcac agtctacatg gagct

35

<210> 123

<211> 35

<212> DNA

<213> Artificial Sequence

<220> Primer

00222E0" 86060550

<223> Synthetic DNA

<400> 123

agctccatgt agactgtgct cgaggacgtg tctgc

35

<210> 124

<211> 26

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 124

agagtcacca tcaccgcaga caagtc

26

<210> 125

<211> 26

<212> DNA

<213> Artificial Sequence

<220> Primer

<223> Synthetic DNA

<400> 125

gacttgtctg cggtgatggt gactct

26

<210> 126

<211> 418

<212> DNA

<213> Artificial Sequence

<220> DNA coding for humanized H chain V region(version s)  
of HM1.24 antibody

<223>

<400> 126  
 atg gac tgg acc tgg agg gtc ttc ttc ttg ctg gct gta gct cca ggt 48  
 Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly  
 -15 -10 -5  
 gct cac tcc cag gtg cag ctg gtg cag tct ggg gct gag gtg aag aag 96  
 Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys  
 -1 1 5 10  
 cct ggg gcc tca gtg aag gtt tcc tgc aag gca tct gga tac acc ttc 144  
 Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe  
 15 20 25  
 act ccc tac tgg atg cag tgg gtg cga cag gcc cct gga caa ggg ctt 192  
 Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu  
 30 35 40 45  
 gag tgg atg gga tct att ttt cct gga gat ggt gat act agg tac agt 240  
 Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser  
 50 55 60  
 cag aag ttc aag ggc aga gtc acc atc acc gca gac aag tcc acg agc 288  
 Gln Lys Phe Lys Gly Arg Val Thr Ile Thr Ala Asp Lys Ser Thr Ser  
 65 70 75  
 aca gcc tac atg gag ctg agc agc ctg aga tct gag gac acg gcc gtg 336  
 Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val  
 80 85 90  
 tat tac tgt gcg aga gga tta cga cga ggg ggg tac tac ttt gac tac 384  
 Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr  
 95 100 105  
 tgg ggg caa ggg acc acg gtc acc gtc tcc tca g 418  
 Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser  
 110 115 120

<210> 127  
 <211> 139  
 <212> PRT  
 <213> Artificial Sequence

<220> Humanized H chain V region(version s) of anti-HM1.24  
 antibody

<223>

002220" 85050550

<400> 127

```

Met Asp Trp Thr Trp Arg Val Phe Phe Leu Leu Ala Val Ala Pro Gly
      -15      -10      -5
Ala His Ser Gln Val Gln Leu Val Gln Ser Gly Ala Glu Val Lys Lys
      -1  1      5      10
Pro Gly Ala Ser Val Lys Val Ser Cys Lys Ala Ser Gly Tyr Thr Phe
      15      20      25
Thr Pro Tyr Trp Met Gln Trp Val Arg Gln Ala Pro Gly Gln Gly Leu
      30      35      40      45
Glu Trp Met Gly Ser Ile Phe Pro Gly Asp Gly Asp Thr Arg Tyr Ser
      50      55      60
Gln Lys Phe Lys Gly Arg Val Thr Ile Thr Ala Asp Lys Ser Thr Ser
      65      70      75
Thr Ala Tyr Met Glu Leu Ser Ser Leu Arg Ser Glu Asp Thr Ala Val
      80      85      90
Tyr Tyr Cys Ala Arg Gly Leu Arg Arg Gly Gly Tyr Tyr Phe Asp Tyr
      95      100      105
Trp Gly Gln Gly Thr Thr Val Thr Val Ser Ser
110      115      120

```

<210> 128

<211> 1013

<212> DNA

<213> Human

<220> DNA coding for HM1.24 antigenic protein

<223>

<400> 128

```

gaattcggca cgagggatct gg atg gca tct act tcg tat gac tat tgc      49
      Met Ala Ser Thr Ser Tyr Asp Tyr Cys
      1      5
aga gtg ccc atg gaa gac ggg gat aag cgc tgt aag ctt ctg ctg ggg      97
Arg Val Pro Met Glu Asp Gly Asp Lys Arg Cys Lys Leu Leu Leu Gly
      10      15      20      25
ata gga att ctg gtg ctc ctg atc atc gtg att ctg ggg gtg ccc ttg      145
Ile Gly Ile Leu Val Leu Leu Ile Ile Val Ile Leu Gly Val Pro Leu
      30      35      40

```

att atc ttc acc atc aag gcc aac agc gag gcc tgc cgg gac ggc ctt	193
Ile Ile Phe Thr Ile Lys Ala Asn Ser Glu Ala Cys Arg Asp Gly Leu	
45 50 55	
cgg gca gtg atg gag tgt cgc aat gtc acc cat ctc ctg caa caa gag	241
Arg Ala Val Met Glu Cys Arg Asn Val Thr His Leu Leu Gln Gln Glu	
60 65 70	
ctg acc gag gcc cag aag ggc ttt cag gat gtg gag gcc cag gcc gcc	289
Leu Thr Glu Ala Gln Lys Gly Phe Gln Asp Val Glu Ala Gln Ala Ala	
75 80 85	
acc tgc aac cac act gtg atg gcc cta atg gct tcc ctg gat gca gag	337
Thr Cys Asn His Thr Val Met Ala Leu Met Ala Ser Leu Asp Ala Glu	
90 95 100 105	
aag gcc caa gga caa aag aaa gtg gag gag ctt gag gga gag atc act	385
Lys Ala Gln Gly Gln Lys Lys Val Glu Glu Leu Glu Gly Glu Ile Thr	
110 115 120	
aca tta aac cat aag ctt cag gac gcg tct gca gag gtg gag cga ctg	433
Thr Leu Asn His Lys Leu Gln Asp Ala Ser Ala Glu Val Glu Arg Leu	
125 130 135	
aga aga gaa aac cag gtc tta agc gtg aga atc gcg gac aag aag tac	481
Arg Arg Glu Asn Gln Val Leu Ser Val Arg Ile Ala Asp Lys Lys Tyr	
140 145 150	
tac ccc agc tcc cag gac tcc agc tcc gct gcg gcg ccc cag ctg ctg	529
Tyr Pro Ser Ser Gln Asp Ser Ser Ser Ala Ala Ala Pro Gln Leu Leu	
155 160 165	
att gtg ctg ctg ggc ctc agc gct ctg ctg cag tga gatccaggga	575
Ile Val Leu Leu Gly Leu Ser Ala Leu Leu Gln ***	
170 175 180	
agctggcaca tcttggaagg tccgtcctgc toggcttttc gcttgaacat tcccttgatc	635
tcatacagttc tgagcgggtc atggggcaac acggttagcg gggagagcac ggggtagccg	695
gagaaggggc tctggagcag gtctggaggg gccatggggc agtcctgggt ctggggacac	755
agtogggttg acccagggct gtctccctcc agagcctccc tccggacaat gagtcccccc	815
tcttgctctcc caccctgaga ttgggcatgg ggtgcggtgt ggggggcatg tgctgcctgt	875
tgttatgggt tttttttgcg ggggggggtt cttttttctg gggtctttga gctccaaaaa	935
aataaacact tcccttgagg gagagcacac cttaaaaaaa aaaaaaaaaa aaaaaaaaaa	995
aaaattoggg cggccgcc	1013

&lt;210&gt; 129

&lt;211&gt; 180

<212> PRT

<213> Human

<220> HM1.24 antigenic protein

<223>

<400> 129

Met Ala Ser Thr Ser Tyr Asp Tyr Cys Arg Val Pro Met Glu Asp Gly  
1 5 10 15  
Asp Lys Arg Cys Lys Leu Leu Leu Gly Ile Gly Ile Leu Val Leu Leu  
20 25 30  
Ile Ile Val Ile Leu Gly Val Pro Leu Ile Ile Phe Thr Ile Lys Ala  
35 40 45  
Asn Ser Glu Ala Cys Arg Asp Gly Leu Arg Ala Val Met Glu Cys Arg  
50 55 60  
Asn Val Thr His Leu Leu Gln Gln Glu Leu Thr Glu Ala Gln Lys Gly  
65 70 75 80  
Phe Gln Asp Val Glu Ala Gln Ala Ala Thr Cys Asn His Thr Val Met  
85 90 95  
Ala Leu Met Ala Ser Leu Asp Ala Glu Lys Ala Gln Gly Gln Lys Lys  
100 105 110  
Val Glu Glu Leu Glu Gly Glu Ile Thr Thr Leu Asn His Lys Leu Gln  
115 120 125  
Asp Ala Ser Ala Glu Val Glu Arg Leu Arg Arg Glu Asn Gln Val Leu  
130 135 140  
Ser Val Arg Ile Ala Asp Lys Lys Tyr Tyr Pro Ser Ser Gln Asp Ser  
145 150 155 160  
Ser Ser Ala Ala Ala Pro Gln Leu Leu Ile Val Leu Leu Gly Leu Ser  
165 170 175  
Ala Leu Leu Gln  
180.